



# Perioperative care in open aortic vascular surgery: A consensus statement by the Enhanced Recovery After Surgery (ERAS) Society and Society for Vascular Surgery

Katharine L. McGinagle, MD, MPH,<sup>a</sup> Emily L. Spangler, MD, MS,<sup>b</sup> Adam C. Pichel, MB, ChB, FRCA,<sup>c</sup> Katie Ayyash, MBChB, BSc, FRCA, MSc,<sup>d</sup> Shipra Arya, MD, MS,<sup>e</sup> Alberto M. Settembrini, MD,<sup>f</sup> Joy Garg, MD,<sup>g</sup> Merin M. Thomas, PA-C,<sup>h</sup> Kate E. Dell, DNP, AGACNP-BC,<sup>i</sup> Iris J. Swiderski, DHSc, MPAS, PA-C,<sup>j</sup> Fae Lindo, NP,<sup>k</sup> Mark G. Davies, MD, PhD, MBA,<sup>l</sup> Carlo Setacci, MD,<sup>m</sup> Richard D. Urman, MD, MBA,<sup>n</sup> Simon J. Howell, MA, MRCP, FRCA, MSc, MD,<sup>o</sup> Olle Ljungqvist, MD, PhD,<sup>p</sup> and Hans D. de Boer, MD, PhD, BC,<sup>q</sup> Chapel Hill, NC; Birmingham, AL; Manchester, York, and Leeds, UK; Palo Alto, CA; Milan, Italy; New Hyde Park, NY; Lafayette, IN; Melbourne, FL; San Antonio, TX; Siena, Italy; Boston, MA; Orebro, Sweden; and Groningen, The Netherlands

## ABSTRACT

The Society for Vascular Surgery and the Enhanced Recovery After Surgery Society formally collaborated and elected an international, multidisciplinary panel of experts to review the literature and provide evidence-based recommendations related to all the health care received in the perioperative period for patients undergoing open abdominal aortic operations (both transabdominal and retroperitoneal approaches, including supraceliac, suprarenal, and infrarenal clamp sites) for aortic aneurysm and aortoiliac occlusive disease. Structured around the Enhanced Recovery After Surgery core elements, 36 recommendations were made and organized into preadmission, preoperative, intraoperative, and postoperative recommendations. (J Vasc Surg 2022;75:1796-820.)

**Keywords:** Abdominal aortic aneurysm; Analgesia and anesthesia; Aortic occlusive disease; Evidence-based recommendations; Enhanced recovery after surgery; Guidelines; Perioperative care

## TABLE OF CONTENTS

<b>DISCLAIMER</b> .....	1797
<b>METHODS</b> .....	1798
Formation of the guideline development group and selection of guideline topics .....	1798
Literature search strategy .....	1798
Quality assessment, data analyses and consensus generation .....	1798

<b>RESULTS</b> .....	1798
<b>PREADMISSION RECOMMENDATIONS</b> .....	1798
1. Patient information, education, and counseling .....	1798
2. Preadmission screening, assessment, and optimization (a, anemia screening; b, nutritional deficiency; c, frailty; d, delirium risk; e, tobacco and alcohol overconsumption; f, medical risk) .....	1799

From the Department of Surgery, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill<sup>a</sup>; the Department of Surgery, School of Medicine, University of Alabama at Birmingham, Birmingham<sup>b</sup>; the Department of Anaesthesia, Manchester Royal Infirmary, Manchester University NHS Foundation Trust, Manchester<sup>c</sup>; the Department of Perioperative Medicine (Merit), York and Scarborough Teaching Hospitals NHS Foundation Trust, York<sup>d</sup>; the Department of Surgery, School of Medicine, Stanford University, Palo Alto<sup>e</sup>; the Fondazione IRCCS, Cà Granda Policlinico, Milan<sup>f</sup>; Department of Vascular Surgery, Kaiser Permanente San Leandro, San Leandro<sup>g</sup>; the Lenox Hill Hospital, Northwell Health, New Hyde Park<sup>h</sup>; the Ascension Medical Group, Lafayette<sup>i</sup>; the Medical Associates of Brevard, Melbourne<sup>j</sup>; the Stanford University Hospital, Palo Alto<sup>k</sup>; the Department of Surgery, Joe R. & Teresa Lozano Long School of Medicine, University of Texas Health Sciences Center, San Antonio<sup>l</sup>; the Department of Surgery, University of Siena, Siena<sup>m</sup>; the Department of Anesthesiology, Perioperative and Pain Medicine, Brigham and Women's Hospital, Boston<sup>n</sup>; the Leeds Institute of Medical Research at St. James's, University of Leeds, Leeds<sup>o</sup>; the Department of Surgery, School of Medical Sciences, Orebro University, Orebro<sup>p</sup>; and the Department of Anesthesiology, Pain Medicine and Procedure Sedation and Analgesia, Martini General Hospital Groningen, Groningen<sup>q</sup>

Author conflict of interest: H.D.B. reports ERAS Society Executive Committee; Global Advisory Board of Merck & Co.; Scientific Advisory Board of Senzime; Global Advisory Board of NMD Pharma; Research and Scientific Committees of the European Society of Anaesthesiology and Intensive Care; Receiving research grants from Merck & Co, and The Medicines Company; S.J.H. reports

Consultancy fees from Edwards Life Sciences and Research funding from Abbott; O.L. reports co-founder ERAS Society, founder and stock owner of Encare AB, advisor Nutricia A/S (The Netherlands), Advisor Abbott (USA), Advisor Merck (USA), Advisor AMN (Canada), speaker honoraria from Fesenius-Kabi, Braun, Nutricia, Nestle, Merck, Encare, Medtronic, previous patent holder of carbohydrate drink; K.L.M. reports speaking honorarium from Medtronic, Inc; R.D.U. reports Fees and/or funding: Pfizer, Merck, Covidien, AcclRx, Heron, Cook; Federal funding: NIH, NSF; Vice-President of ERAS USA. E.L.S., A.C.P., K.A., S.A., A.M.S., M.M.T., K.E.D., I.J.S., F.L., M.G.D., C.S. have no conflicts of interest.

Correspondence: Katharine L. McGinagle, MD, MPH, Department of Surgery, School of Medicine, University of North Carolina at Chapel Hill, 3022 Burnett-Womack Bldg, CB 7212, Chapel Hill, NC 27599 (e-mail: [katharine\\_mcginagle@med.unc.edu](mailto:katharine_mcginagle@med.unc.edu)).

Independent peer review and oversight have been provided by the members of the Society for Vascular Surgery Document Oversight Committee (Ruth Bush, chair, Marc Schermerhorn, vice-chair, Keith Calligaro, Mohammad Eslami, Raul Guzman, Ahmed Kayssi, Chris Kwolek, Gregory Landry, Erika Mitchell, Patrick Muck, William Robinson, Jeannie Ruddy, Evan Ryer, Palma Shaw, Chris Smolock, Britt Tonnessen, Brant Ulley, Chandu Vemuri, Greg Westin, and Karen Woo).

0741-5214

Copyright © 2022 Published by Elsevier Inc. on behalf of the Society for Vascular Surgery.

<https://doi.org/10.1016/j.jvs.2022.01.131>

2a. Anemia screening .....	1799
2b. Nutritional deficiency .....	1799
2c. Frailty .....	1799
2d. Delirium risk .....	1802
2e. Tobacco and alcohol cessation .....	1802
2f. Medical risk .....	1803
2f-i. Cardiovascular risk .....	1803
2f-ii. Renal risk .....	1804
3. Preoperative exercise therapy and prehabilitation .....	1804
4. Perioperative anticoagulation strategy or plan .....	1804
<b>PREOPERATIVE RECOMMENDATIONS .....</b>	<b>1805</b>
5. Fasting .....	1805
6. Carbohydrate loading .....	1805
7. Venous thromboembolism prophylaxis .....	1805
8. Preanesthetic sedative and analgesia medication .....	1806
9. Antimicrobial prophylaxis and skin preparation .....	1806
10. Prevention of nausea and vomiting .....	1806
<b>INTRAOPERATIVE RECOMMENDATIONS .....</b>	<b>1807</b>
11. Standard anesthetic protocol .....	1807
11a. Neuromuscular blocking agents and monitoring .....	1807
11b. Cardiovascular monitoring .....	1807
11c. Renal protection .....	1808
11c-i. Ischemic preconditioning .....	1808
11c-ii. Renal protective medications .....	1808
11d. Cell salvage .....	1808
11e. Lung protective ventilation .....	1808
11f. Epidural analgesia .....	1809
12. Body temperature management .....	1809
13. Drainage of surgical site strategy .....	1809
<b>POSTOPERATIVE RECOMMENDATIONS .....</b>	<b>1809</b>
14. Multimodal analgesia and opioid reduction .....	1809
15. Nasogastric drainage .....	1810
16. Oral feeding .....	1810
17. Fluid therapy .....	1810
18. Urinary drainage .....	1810
19. Glycemic control .....	1811
20. Early mobilization .....	1811
21. Discharge education .....	1811
22. Audit of outcomes .....	1812
<b>CARE ELEMENTS WITH CONFLICTING OR INSUFFICIENT DATA FOR A RECOMMENDATION .</b>	<b>1812</b>
Preoperative medical risk assessment .....	1812
Skin preparation .....	1812
Intraoperative fluid therapy .....	1813
<b>SUMMARY AND CONCLUSIONS .....</b>	<b>1813</b>

## DISCLAIMER

The Society for Vascular Surgery (SVS) develops evidenced-based documents as a resource to assist members in the practice of vascular surgery. The recommendations contained herein are based on a recent review of published evidence and expert opinion. They reflect the available body of evidence, and their

applicability reflects the limitations of those data and are subject to reassessment and revision as new knowledge emerges. Given these limitations, consensus documents do not represent a statement of the standard of care, nor do they substitute for clinician judgment or supplant patient preference or shared decision-making. The Society for Vascular Surgery recognizes that departure from these recommendations could be warranted when, in the reasonable judgment of the treating clinician, such a course of action is indicated by the clinical presentation of the patient, limitations of available resources, advances in knowledge or technology, or patient preference. The reader must rely solely on their own judgment to determine which practices and procedures, whether included in this document or not, are appropriate for them, their patient, their institution, or their practice.

Enhanced Recovery After Surgery (ERAS) pathways have been shown to be beneficial for many surgical specialties.<sup>1</sup> The concept of multidisciplinary care to control postoperative physiology and maintain homeostasis to attenuate stress and accelerate convalescence was first described in 1997.<sup>2</sup> Soon thereafter, the perioperative enhanced recovery after surgery study group was formed in Europe, and their first consensus statement for perioperative care to enhance recovery after colorectal surgery was published.<sup>3</sup> The group formed the ERAS Society in 2010. Its mission has since spread to six continents, and 19 surgical specialties have published formal ERAS Society guidelines or consensus statements (available at: [www.erassociety.org](http://www.erassociety.org)). Although each specialty and geographic region might be unique, the common foundational theme across all ERAS guidelines and consensus statements is to create a culture in which all perioperative caregivers, including anesthesiologists, surgeons, nurses, and therapists work together within a clinical pathway to apply evidence-based practice in unison. In addition to these recommendations, auditing systems with data points corresponding to each ERAS element are supported by the ERAS Society, allowing those who choose to participate a method to monitor the fidelity of their enhanced recovery programs.

Compared with the general population, vascular surgery patients are more likely to present with specific perioperative challenges owing to age, frailty, and multiple comorbidities and the highly invasive operations that are sometimes required for limb or life preservation. This combination of factors results in complex management strategies, increased usage of health care resources, prolonged hospitalization stays, and increased rehabilitation needs. ERAS, with its aim of delivering high-quality perioperative care and accelerating recovery, appears well suited to address the needs of patients undergoing open aortic operations. The present

consensus statement focused on both transabdominal and retroperitoneal approaches (including supraceliac, suprarenal, and infrarenal clamp sites) for aortic aneurysm and aortoiliac occlusive disease operations.

## METHODS

**Formation of the guideline development group and selection of guideline topics.** In collaboration, the SVS and the ERAS Society initiated the formation of the multidisciplinary, international guideline development group (GDG) comprising vascular surgeons, anesthesiologists, and advanced practice providers with expertise in ERAS and vascular pathophysiology. Guideline topic selection was performed according to the recommendations for the development of clinical guidelines within the ERAS Society framework.<sup>4</sup> The GDG was consulted for advice on the appropriate items to be included in the guideline, with the final decision made by the lead authors (K.L.M. and H.D.B.). The guideline items were allocated to the authors for literature summary and grading, depending on each individual's expertise. The final content was agreed on by all the authors.

**Literature search strategy.** A third-party team of investigators and a reference librarian specializing in literature reviews assisted the GDG in identifying and categorizing the existing effectiveness and comparative effectiveness literature on ERAS protocols relevant to vascular surgery. The librarian executed and documented the search in July 2019. The databases queried included PubMed, CINAHL, and [ClinicalTrials.gov](https://www.clinicaltrials.gov). Systematic reviews, randomized controlled trials (RCTs), and observational cohort studies reporting on adults (age  $\geq 18$  years) undergoing aortic surgery and reporting on any one or combinations of the ERAS guideline items were included. Many items had not been studied for open aortic surgery patients; thus, for elements that could be reasonably applied to all types of surgical patients, the reported data and ERAS guidelines from other surgical specialties were included. In some cases, the data from aortic and other patient populations were insufficient, or the existing data could not be reasonably extrapolated. Thus, no recommendations were made. The GDG members then searched reference lists and also repeated literature searches to include any new publications up to and including April 2021. The final included studies were carefully reviewed by the GDG, and any disagreements were resolved through group consensus via the Delphi method.

**Quality assessment, data analyses and consensus generation.** The GRADE (grading of recommendations, assessment, development, and evaluation) system was used to evaluate the quality of evidence and recommendations for each of the ERAS topics.<sup>5,6</sup> Recommendations

were made according to whether the quality of evidence was high, moderate, low, or very low. Conclusions based on high-quality evidence are unlikely to change with further investigation. Those based on moderate-quality evidence are more likely to be affected by future research. Finally, those based on low-quality evidence are the least supported, might not include prospective study designs, and are the most likely change in the future if more rigorous research is performed. In the case of recommendations made solely from literature of nonaortic cohorts, the quality of evidence was downgraded. The strength of the recommendation was based on the balance between the desirable and undesirable effects of the recommendation. A strong recommendation for an ERAS item is possible even with a low quality of evidence, if the risk of harm is negligible.

## RESULTS

The literature search yielded 3460 citations. Of these 3460 citations, 109 studies had addressed enhanced recovery interventions, as either individual core elements or combinations of elements, for vascular surgery patients. Another 18 studies had included mixed cohorts, including vascular surgery patients. In addition, 463 studies were identified that had not included vascular surgery patients but that might have been applicable to patients undergoing open aortic operations. A total of 590 of these abstracts were screened. After discussions and reviews, no disagreements were found between the authors in the assessment of the quality of evidence and grading of the included reports. Structured around the ERAS core elements, 36 recommendations were made and organized into preadmission, preoperative, intraoperative, and postoperative recommendations (Table; Fig).<sup>7</sup> For some of the ERAS elements, insufficient data were found to support a recommendation; however, summary statements based on expert opinion were provided.

## PREADMISSION RECOMMENDATIONS

### 1. Patient information, education, and counseling

Preoperative education and counseling are essential to engage patients and caregivers in the perioperative process. A recent systematic review and meta-analysis showed that two thirds of studies (two RCTs and five observational studies) of different enhanced recovery programs for open aortic surgeries had used some form of patient education, although education had not been studied as a specific intervention.<sup>8</sup> A retrospective study showed effectiveness in enhanced recovery after open abdominal aortic aneurysm (AAA) repair through counseling before and during hospitalization regarding nutrition, early rehabilitation, and setting expectations and daily goals.<sup>9</sup> Preoperative education can be provided using various methods, such as illustrated booklets,<sup>10</sup> in addition to written informed consent, surgeon patient discussion,<sup>11</sup> and

multidisciplinary education with surgery and anesthesia services.<sup>12</sup> Counseling of patients and families regarding early mobilization<sup>10,11</sup> and discharge criteria<sup>13</sup> are essential to achieve length of stay (LOS) reductions. Studies, including an RCT, from other surgical specialties have shown that extended pre- and postoperative counseling, guidance by an ERAS nurse, and written information will lead to less pain, decreased nausea and vomiting, reduced hospital LOS, and increased enhanced recovery pathway adherence.<sup>14-16</sup>

*Recommendation:* Patients should receive dedicated verbal and written preoperative education and counseling.

*Quality of evidence:* C (low)

*Strength of recommendation:* Grade 1 (strong)

## 2. Preadmission screening, assessment, and optimization (a, anemia screening; b, nutritional deficiency; c, frailty; d, delirium risk; e, tobacco and alcohol overconsumption; f, medical risk)

Delivery of optimal care in aneurysm disease requires balancing the operative risk relative to the likelihood of survival in the perioperative period and beyond. A recent systematic review and meta-analysis of RCTs and observational studies found that only 3 of 12 studies of ERAS-like pathways in aortic surgery had had any medical screening and preoperative optimization documented.<sup>8,9,11,17</sup> Patients should undergo a thorough preoperative assessment and optimization across a breadth of risk factors for aortic surgery, as detailed in the next sections.

**2a. Anemia screening.** Anemia is associated with poor outcomes after surgery.<sup>18,19</sup> Although a preoperative blood transfusion will correct anemia rapidly and can be used for severely anemic patients and/or patients undergoing surgery with expected profound blood loss, caution should be used because transfusions have been associated with increased mortality and morbidity.<sup>20,21</sup> Medical management of preoperative anemia requires time and should be planned  $\geq 3$  to 4 weeks before elective surgery. However, aortic surgery-specific evidence has remained lacking. Extrapolated from other specialties, iron (oral and intravenous), folate, and vitamin B<sub>12</sub> supplements might reduce the need for blood transfusions but did not affect the incidence of adverse events or mortality.<sup>22-27</sup> In colorectal ERAS pathways, iron supplementation has been recommended.<sup>28</sup> However, the balance of trial evidence in mixed surgical populations has not shown clinically significant reductions in the need for allogeneic blood transfusions for patients receiving preoperative iron therapy.<sup>29,30</sup> Caution is advised in translating certain anemia therapies to open aortic operations amid

the presence of U.S. Food and Drug Administration limitations on the use of erythropoietin-stimulating agents for patients undergoing cardiac or vascular surgery and an adverse event profile that includes thrombotic events.<sup>31</sup>

*Recommendation:* The cause of preoperative anemia should be evaluated and treated.

*Quality of evidence:* D (very low)

*Strength of recommendation:* Grade 2 (weak)

**2b. Nutritional deficiency.** Preoperative nutritional deficiency can be established by screening and, as needed, assessed using a combination of clinical and biochemical parameters. Several assessment tools are available to assist in this regard, including Malnutrition Universal Screening Tool, Nutritional Risk Screening 2002, Mini Nutritional Assessment, Short Nutritional Assessment Questionnaire, Malnutrition Screening Tool, and Subjective Global Assessment.<sup>32-34</sup> In a single-site retrospective study of open AAA repairs, a moderate risk of malnutrition determined from Controlling Nutritional Status scores (derived from serum albumin, cholesterol levels, and lymphocyte counts in peripheral blood) was associated with mid-term mortality and life-threatening complications.<sup>35</sup>

If the screening results indicate risk, a nutritional assessment should be performed if expertise is available and appropriate nutritional care delivered.<sup>36,37</sup> Oral nutritional supplements, including those containing immunonutrients,<sup>38-40</sup> should almost always be preferred to enteral or parenteral nutrition regimens.<sup>37</sup> Regimens for correcting malnutrition via oral nutritional supplementation have shown the greatest effect if started  $\geq 7$  to 10 days preoperatively in nonvascular cohorts.<sup>32,41-43</sup>

*Recommendation:* Screening should be performed for malnutrition and nutritional deficiencies corrected, preferably with oral regimens.

*Quality of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

**2c. Frailty.** Frailty is a syndrome defined as increased vulnerability because of a physiologic decline in reserve and function and consists of physical, functional, social, nutritional, and cognitive domains. For aortic aneurysm patients, frailty has been associated with increased adverse outcomes at 30 days (ie, morbidity, mortality, failure to rescue) and an increased LOS.<sup>44,45</sup> Preoperative frailty is associated with increased discharge to non-home destination, a loss of independence after major vascular surgery, and decreased long-term survival.<sup>46-48</sup> Often used scales to measure frailty include the Clinical Frailty Scale, modified Frailty index, and the Risk Analysis Index. The use of standardized frailty assessments for aneurysm patients more than doubles the pool of the at-risk or frail patients identified for preoperative optimization.<sup>49</sup>

**Table.** Recommendations and GRADE (grading of recommendations, assessment, development, evaluation) for each Enhanced Recovery After Surgery (ERAS) element

ERAS element	Recommendation	GRADE	Evidence level
Preadmission			
1. Patient information, education, counseling	Patients should receive dedicated verbal and written preoperative education and counseling (grade 1C)	Strong	Low <sup>a</sup>
2a. Screening, assessment, and optimization: anemia screening	Evaluate the cause of and treat chronic preoperative anemia (grade 2D)	Weak	Very low <sup>a</sup>
2b. Screening, assessment, and optimization: nutritional deficiency	Screen for malnutrition and correct nutritional deficiency, preferably with oral regimens (grade 1B)	Strong	Moderate <sup>a</sup>
2c. Screening, assessment, and optimization: frailty	Frailty screening is recommended as a routine part of preoperative patient assessment (grade 1B)	Strong	Moderate
2d. Screening, assessment, and optimization: delirium risk	Screen for delirium risk and implement preoperative practices to minimize onset of delirium as a routine part of practice (grade 1C)	Strong	Low
2e. Screening, assessment, and optimization: tobacco and alcohol			
	i. Prescribe smoking cessation therapy and recommend a minimum of 4 weeks smoking cessation before surgery (grade 1A)	Strong	High <sup>a</sup>
	ii. Recommend alcohol cessation 4 weeks before surgery, particularly for patients who consume more than five alcohol equivalents or 15 alcohol units per day (grade 1C)	Strong	Low <sup>a</sup>
2f. Screening, assessment, and optimization: medical risk			
	i. Cardiac risk should be evaluated and optimized with antiplatelet therapy, statins, antihypertensive therapy, and diabetes control before elective surgery (grade 1A)	Strong	High
	ii. Risk of acute kidney injury should be evaluated and optimized before elective surgery (grade 1A)	Strong	High
3. Preoperative exercise therapy and rehabilitation	Recommend 6 weeks of supervised exercise therapy before elective surgery (grade 2C)	Weak	Low
4. Perioperative antiplatelet or anticoagulation plan			
	i. Continue aspirin throughout the perioperative period (grade 1A)	Strong	High
	ii. Stop secondary antiplatelet medications and anticoagulant agents before surgery, except for select patient populations (grade 1B)	Strong	Moderate
Preoperative			
5. Fasting	Avoid overnight fasting; encourage clear fluids for $\leq 2$ hours and light foods for $\leq 6$ hours before the induction of general anesthesia (grade 1A)	Strong	High <sup>a</sup>
6. Carbohydrate loading	Patients without diabetes should receive a preoperative carbohydrate drink (grade 2B)	Weak	Moderate <sup>a</sup>
7. Venous thromboembolism prophylaxis	Routinely use calf-length intermittent compression devices combined with either low-dose unfractionated heparin or low-molecular-weight heparin starting immediately before surgery and continuing at least throughout the hospitalization (grade 1B)	Strong	Moderate <sup>a</sup>
8. Preanesthetic sedative and analgesia medication			
	i. Do not routinely use sedatives to reduce anxiety preoperatively (grade 1C)	Strong	Low <sup>a</sup>

**Table.** Continued.

ERAS element	Recommendation	GRADE	Evidence level
	ii. Routinely use preoperative administration of acetaminophen, NSAIDs, and gabapentinoids as part of a multimodal opioid-sparing analgesia strategy (grade 1C)	Strong	Low <sup>a</sup>
9. Antimicrobial agents	Prophylactic intravenous antibiotic dosing should begin 30-60 minutes preoperatively, with redosing intraoperatively within two serum half-lives of the antimicrobial agent used or with substantive intraoperative blood loss, and extending no more than 24 hours postoperatively (grade 1A)	Strong	High
10. Prevention of nausea and vomiting	Perform a risk assessment for PONV, routinely use multimodal PONV prophylaxis based on the assessment findings, and use PONV rescue with a different class of antiemetic (grade 1A)	Strong	High <sup>a</sup>
Intraoperative			
11a. Anesthetic protocol: neuromuscular blocking agents and monitoring	Neuromuscular monitoring should be used to ensure adequate intraoperative muscle relaxation and full reversal of neuromuscular blockade before extubation (grade 1A)	Strong	High <sup>a</sup>
11b. Anesthetic protocol: cardiovascular monitoring	Invasive cardiovascular monitoring with an arterial catheter is essential; the MAP should be maintained to near baseline values and at or >65 mm Hg (grade 1B)	Strong	Moderate
11c. Anesthetic protocol: renal protection	Adjunctive renal protective medications and methods of ischemic preconditioning of any type can cause harm and should not be used for infrarenal repairs (grade 1B)	Strong	Moderate
11d. Anesthetic protocol: cell salvage	Use cell salvage (grade 1B)	Strong	Moderate
11e. Anesthetic protocol: lung-protective ventilation	Use lung protective strategies (grade 2C)	Weak	Low <sup>a</sup>
11f. Epidural analgesia	Mid-thoracic (T6-T9) epidural analgesia should be used intraoperatively and continued postoperatively as an infusion or patient-controlled analgesia using a combination of a local anesthetic and an opioid (grade 1B)	Strong	Moderate
12. Body temperature management	Use multiple strategies to maintain normothermia, including prewarming and active warming of patients intraoperatively (grade 1B)	Strong	Moderate <sup>a</sup>
13. Drainage of surgical site	Do not routinely use drains in the surgical wound (grade 2C)	Weak	Low
Postoperative			
14. Multimodal analgesia and opioid reduction strategies	Routinely use multimodal analgesic regimens to improve pain control and reduce opioid consumption (grade 1B)	Strong	Moderate <sup>a</sup>
15. Nasogastric drainage	Avoid the routine use of postoperative nasogastric tubes (grade 1A)	Strong	High <sup>a</sup>
16. Oral feeding	An early return to a normal diet and oral nutritional supplements should be promoted (grade 1B)	Strong	Moderate
17. Fluid therapy	If intravenous fluids are indicated, restrict postoperative fluid therapy to 1.5 L/d (grade 1B)	Strong	Moderate
18. Urinary drainage	Early removal of urinary drainage catheters is recommended (grade 1C)	Strong	Low <sup>a</sup>
19. Glycemic control	Maintain glycemic control, with attention to avoiding hypoglycemia (grade 1B)	Strong	Moderate

(Continued on next page)

Table. Continued.

ERAS element	Recommendation	GRADE	Evidence level
20. Early mobilization strategy	Use a formal plan for early mobilization, with early physical therapy involvement when needed (grade 1C)	Strong	Low <sup>a</sup>
21. Discharge education	Use both written and verbal discharge education and follow-up telephone calls within 24-48 hours of hospital discharge and arrange clinic follow-up visits with both the surgeon and the primary care physician at least once by 30 days postoperatively (grade 1C)	Strong	Low <sup>a</sup>
22. Audit of outcomes	Routine auditing of ERAS care processes and feedback to all staff involved are necessary (grade 1C)	Strong	Low <sup>a</sup>

MAP, Mean arterial pressure; NSAIDs, nonsteroidal anti-inflammatory drugs; PONV, postoperative nausea and vomiting.  
<sup>a</sup>Some of the evidence used to make the recommendation was extrapolated from nonaortic patient cohorts.

Based on the consensus of best practices from the Society for Perioperative Assessment and Quality Improvement, a patient who has a positive frailty screening result should be followed up with a diagnostic assessment of frailty, and, when feasible, a comprehensive geriatric assessment with a tailored intervention (shared decision-making or prehabilitation) should be performed, ideally by a geriatric specialist.<sup>50</sup> Specifically, a small RCT testing an outpatient comprehensive geriatric assessment and optimization program for elderly patients scheduled for elective vascular surgery was associated with reduced complications, shorter LOSs, and a lower rate of patients discharged to a higher level of care dependency.<sup>51</sup>

*Recommendation:* Frailty screening is recommended as a routine part of preoperative patient assessments.

*Quality of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

**2d. Delirium risk.** Delirium can contribute to adverse events, prolonged hospital LOSs, non-home discharge destinations, and mortality following surgery.<sup>52</sup> Those with a medical history of delirium, advancing age by decade, and American Society of Anesthesiologists score of  $\geq 3$  were at the highest risk of delirium in a prospective cohort of patients undergoing surgery for AAAs and colorectal cancer.<sup>52</sup> A subsequent cohort study of the same patient population additionally showed renal impairment, cognitive impairment, active smoking, intensive care unit admission, and blood transfusion were associated with an increased risk of postoperative delirium, but prehabilitation program participation was protective.<sup>53,54</sup>

Preoperative identification of cognitive risk using standardized cognitive scales such as the mini-mental state examination can also allow engagement of patients and/or families to discuss the implications of proposed surgery (capacity to consent and risk of postoperative

complications, including delirium) and the long-term effects of cognitive impairment.<sup>51,55</sup>

*Recommendation:* Patients should undergo screening for delirium risk and preoperative practices to minimize the onset of delirium should be implemented as a routine part of practice.

*Quality of evidence:* C (low)

*Strength of recommendation:* Grade 1 (strong)


**2e. Tobacco and alcohol cessation.** The SVS practice guidelines for AAAs recommend smoking cessation for  $\geq 2$  weeks before aneurysm repair,<sup>56</sup> although the supporting literature is limited. In a systematic review and meta-analysis of 25 studies (2 RCTs, 7 prospective cohort studies, and 16 retrospective studies), including >20,000 cardiac and noncardiac surgical patients, a 23% and 47% reduction was found in the incidence of respiratory complications for patients who had stopped smoking for 4 and 8 weeks before surgery, respectively.<sup>57</sup> Another meta-analysis of four RCTs demonstrated that smoking cessation interventions reduced surgical site infections (SSIs), and pooled data from 140 cohort studies supported the finding that 1 month of smoking cessation was associated with improved postoperative outcomes.<sup>57-59</sup> In the perioperative period, nicotine replacement therapy and smoking cessation medications have proved efficacious for smoking cessation.<sup>60</sup>

*Recommendation:* Smoking cessation therapy should be prescribed and a minimum of 4 weeks smoking cessation before surgery recommended.

*Quality of evidence:* A (high)

*Strength of recommendation:* Grade 1 (strong)

In a Cochrane review of two RCTs comprising 69 surgical patients, overall postoperative morbidity was increased by two- to threefold for patients who had abused alcohol. One month of preoperative abstinence significantly improved the outcomes for a group who



**Fig.** Enhanced Recovery After Surgery (ERAS) diagram graphically showing the timeline for each of the ERAS elements to be performed during the perioperative period. *Intra*, Intraoperative; *PACU*, postanesthesia care unit; *POD*, postoperative day; *Post*, postoperative; *Pre*, preoperative.

had imbibed at least five alcohol equivalents (or 15 units) per day.<sup>61</sup> Another meta-analysis of multiple RCTs in mixed surgical cohorts showed that drinking more than one alcohol equivalent (or two units) per day increased the risk of non-SSI but did not increase postoperative mortality.<sup>62</sup>

*Recommendation:* Alcohol cessation should be recommended for 4 weeks before surgery, especially for patients who consume more than five alcohol equivalents or 15 alcohol units per day.

*Quality of evidence:* C (low)

*Strength of recommendation:* Grade 1 (strong)

**2f. Medical risk.** Delivering optimal care in aortic disease requires balancing the operative risk relative to the likelihood of survival in the perioperative period and beyond. Patients with AAA are known to experience higher rates of heart attack, stroke, major amputation, and 5-year mortality compared with age-matched controls.<sup>63</sup> Patients should undergo a thorough preoperative assessment and optimization across a breadth of risk factors for aortic surgery as detailed in the guidelines for perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery.<sup>64</sup>

Risk scores can be used to quantify the overall risk of open aortic surgery and/or specific risk to a single physiologic system; multiple validated risk calculators are

available.<sup>65-67</sup> The SVS Clinical Practice Guidelines for AAA suggest “informing patients contemplating open repair or endovascular AAA repair of the VQI [Vascular Quality Initiative] perioperative mortality risk score.”<sup>56</sup> The Vascular Quality Initiative mortality risk model predicts for mortality well (C = 0.802).<sup>68</sup> However, the model discrimination for predicting for postoperative myocardial infarction after open aortic surgery was only 0.68.<sup>67</sup> Predicting the risk to a single organ system is less accurate with the available tools, which are not universally recommended.<sup>69</sup>

**2f-i. Cardiovascular risk.** Based on multiple high-quality RCTs of cardiovascular disease prevention, it is recommended that all patients with aneurysmal disease and aortoiliac occlusive disease should be offered antiplatelet therapy, statins (even if the low-density lipoprotein cholesterol level is <100 mg/dL), and antihypertensive agents to maintain the systolic blood pressure at <140 mm Hg.<sup>64,70-72</sup> However, we caution against starting a beta-blocker for blood pressure control immediately before surgery because its use appears to increase the 30-day mortality for noncardiac surgical populations.<sup>73</sup> Diabetes control with oral agents and/or insulin is also recommended because diabetes mellitus is a risk factor for cardiovascular disease, in general, and postoperative major adverse cardiac events, specifically. However, specific hemoglobin A1c targets and associated risks have not been specified in the setting of aortic surgery.<sup>64,74,75</sup>

*Recommendation:* Patients' cardiac risk should be evaluated and optimized with antiplatelet therapy, statins, antihypertensive therapy, and diabetes control before elective surgery.

*Level of evidence:* A (high)

*Strength of recommendation:* Grade 1 (strong)

**2f-ii. Renal risk.** Even in patients with normal baseline renal function, transient postoperative renal impairment has been associated with increased morbidity and the need for critical care support.<sup>76</sup> Furthermore, patients with existing chronic kidney disease face a well-established risk of poor outcomes after open aortic operations.<sup>77</sup> In accordance with the Kidney Disease: Improving Global Outcomes recommendations determined from systematic reviews of RCTs, angiotensin-converting enzyme inhibitors and angiotensin receptor antagonists should be withheld on the morning of surgery and only restarted once euvolemia has been achieved postoperatively.<sup>78</sup> Patients with severe renal insufficiency (ie, chronic kidney disease stage 4 or 5; estimated glomerular filtration rate <30 mL/min/1.73 m<sup>2</sup>) should be evaluated by a specialist to optimize renal function before elective aortic repair.<sup>79</sup>

*Recommendation:* The patient's risk of acute kidney injury should be evaluated and optimized before elective surgery.

*Level of evidence:* A (high)

*Strength of recommendation:* Grade 1 (strong)

### 3. Preoperative exercise therapy and prehabilitation

Although frailty screening can be used in the preoperative discussion about proceeding with surgery, the nascent work in prehabilitation regimens has aimed to reduce the limitations in functional capacity among frail patients to better allow them to withstand the stress of surgery.<sup>80,81</sup> Prehabilitation engages the patient in a combination of exercise training, nutrition therapy, and mental preparation for surgery.<sup>82</sup> Based on several studies within cardiac<sup>83-85</sup> and abdominal surgery,<sup>86-88</sup> 3 to 4 weeks of prehabilitation is considered feasible and has been recommended in other specialties' ERAS guidelines.<sup>89,90</sup>

Studies of the exercise component of prehabilitation have been shown to be beneficial in the recovery after open aortic surgery; however, none of the vascular studies of ERAS-like programs in aortic surgery had had a complete prehabilitation program available.<sup>8</sup> A systematic review of the effects of exercise alone for AAA patients awaiting surgery found only four studies, all of which favored exercise training.<sup>91</sup> One RCT of 124 patients undergoing open or endovascular AAA repair examined the standard treatment vs 6 weeks of preoperative supervised exercise, comprising aerobic, stretching, and resistance training for 1 hour, three times per week.<sup>92</sup>

Overall, a significant decrease was found in postoperative complications and shorter LOSs; however, the study was not powered to analyze open aortic surgery alone. Inspiratory muscle training for 6 d/wk for 2 weeks in 15-minute sessions showed a trend toward a decreased occurrence and duration of atelectasis in the intervention vs control group in another small RCT.<sup>93</sup> A cohort study demonstrated the feasibility of a high intensity interval training regimen for 12 weeks with three sessions per week.<sup>94</sup> Another cohort study found that AAA patients who had exercised early in preparation for surgery compared with later in recovery from surgery had had a significantly shorter LOS and time to attaining independence in sitting, standing, and walking.<sup>95</sup> Although the studies had included a diversity of training programs and had reported mixed results, they favored the use of the exercise component of prehabilitation and showed the safety of exercise for AAA patients with no adverse events related to the aneurysm.

*Recommendation:* Six weeks of supervised exercise therapy before elective surgery should be recommended.

*Level of evidence:* C (low)

*Strength of recommendation:* Grade 2 (weak)

### 4. Perioperative anticoagulation strategy or plan

Based on a RCT and a prospective cohort study of >1600 patients undergoing vascular surgery, aspirin reduces the incidence of adverse cardiovascular events and mortality among patients with coronary artery disease and should be continued during the perioperative period.<sup>96-98</sup> Often, patients with coexistent coronary and aortic disease will be receiving dual antiplatelet therapy, and the 2016 American College of Cardiology/American Heart Association guidelines recommend avoiding discontinuation of the second agent within 1 month after bare metal stent placement and 6 months after drug-eluting coronary stent placement, although the risk of stent thrombosis must be weighed against the risk of delaying aortic surgery on a case-specific basis.<sup>98</sup> These guidelines were determined by from multiple large observational studies of mixed surgical populations evaluating the timeframe for an increased risk of coronary stent thrombosis.<sup>99-102</sup> Practical recommendations for standardizing the perioperative management of antiplatelet and anticoagulation therapy for patients with coronary stents who require vascular surgery have been developed.<sup>103</sup>

When feasible, any antiplatelet medication, other than aspirin, should be stopped 7 days before surgery to reduce the risk of regional anesthesia techniques, including epidural placement, and can be resumed 6 hours after block performance or epidural catheter removal.<sup>104,105</sup> Both warfarin and non-vitamin K

antagonist oral anticoagulants should be discontinued  $\geq 5$  days and 2 days in advance of surgery, respectively, to allow for regional anesthetic techniques and to reduce the risk of intraoperative bleeding.<sup>106</sup> The need for low-molecular-weight heparin as a bridge should be determined by the indication for anticoagulation.<sup>106</sup>

*Recommendation:* Aspirin should be continued throughout the perioperative period.

*Level of evidence:* A (high)

*Strength of recommendation:* Grade 1 (strong)

*Recommendation:* Secondary antiplatelet medications and anticoagulant agents should be stopped before surgery, except for select patient populations.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

## PREOPERATIVE RECOMMENDATIONS

**5. Fasting.** Modern preoperative fasting guidelines have been a part of many enhanced recovery bundles studied in open aortic surgery, and limited fasting is feasible.<sup>8</sup> Based on a meta-analysis of multiple RCTs that included a range of surgical patients and according to both the American Society of Anesthesiologists' and European Society of Anesthesiology's preoperative fasting guidelines, which give strong recommendations based on high-quality evidence, patients should be allowed to drink clear fluids for  $\leq 2$  hours before elective surgery but should not eat any solid food for  $\geq 6$  hours for "light meals" or 8 hours for fatty foods and meat.<sup>107,108</sup> However, it is important to note that these recommendations only apply to patients who will undergo elective procedures, and a longer fasting period might be necessary for patients with delayed gastric emptying.

*Recommendation:* Patients should avoid overnight fasting and should be encouraged to drink clear fluids for  $\leq 2$  hours and light foods  $\leq 6$  hours before the induction of general anesthesia.

*Level of evidence:* A (high)

*Recommendation:* Grade 1 (strong)

**6. Carbohydrate loading.** No RCTs have specifically examined the role of oral carbohydrate loading for patients undergoing open aortic surgery; however, this practice was included as a part of a bundle of care in two "fast-track" aortic surgery publications.<sup>8,9,11</sup> Extrapolated from other specialties, the most commonly studied regimen has been 800 mL of a clear 12.5% carbohydrate drink the evening before surgery and 400 mL  $\leq 2$  hours before surgery.<sup>109,110</sup> Several RCTs of preoperative oral carbohydrate drinks reported that their use can reduce postoperative insulin resistance and result in less perioperative hyperglycemia compared with either placebo or fasting after midnight.<sup>111-113</sup> Other RCTs of variable size and quality have demonstrated that carbohydrate administration can modulate postoperative metabolic

and inflammatory responses, decrease protein breakdown, and better preserve patients' lean body mass and muscle strength compared with preoperative fasting.<sup>114</sup> Several medium-size RCTs of coronary artery bypass surgery have demonstrated that the consumption of preoperative carbohydrates can reduce the incidence of myocardial injury, the intensive care unit LOS, and the need for vasoactive drug administration.<sup>115-117</sup> A systematic review of 22 RCTs found that oral carbohydrate loading is safe for  $\leq 2$  hours before surgery and can improve postoperative discomfort.<sup>118</sup> A Cochrane review of 27 RCTs of variable quality in mixed surgical populations concluded that preoperative carbohydrate loading reduces the hospital LOS, especially after major operations.<sup>119</sup>

Because many patients undergoing vascular surgery will have diabetes, the question of effectiveness and safety has been raised. Despite the low-quality evidence, it appears that oral carbohydrates are safe and effective in patients with diabetes.<sup>120,121</sup> However, no recommendations can be made for patients with diabetes.

*Recommendation:* Patients without diabetes should receive a preoperative carbohydrate drink.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 2 (weak)

**7. Venous thromboembolism prophylaxis.** Based on two RCTs, multiple prospective cohort studies of  $>1500$  patients combined, and a retrospective analysis of registry data, the incidence of deep vein thrombosis (DVT) after open aortic surgery has been high, even among patients receiving perioperative chemoprophylaxis.<sup>122-128</sup> Within the initial hospitalization, the rate of DVT diagnosed with screening venous duplex ultrasound has ranged from 7.5% to 18% and might be as great as 37% within the 30-day postoperative period.<sup>123</sup> Specific to open aortic surgery, one RCT of 100 patients compared unfractionated heparin of 5000 U twice daily and calf-length intermittent mechanical compression to no DVT prophylaxis.<sup>128</sup> They found no differences in the rate of DVT or pulmonary embolism and reported no bleeding events.<sup>128</sup> Another RCT of 224 patients compared the effectiveness of unfractionated heparin to low-molecular-weight heparin and found a high rate of DVT (7.5% within 10 days) but no differences in effectiveness between the treatment groups.<sup>122</sup> Three patients (1%) had developed a wound hematoma requiring exploration; however, no major bleeding events were reported.<sup>122</sup>

Although aortic surgery is considered high risk for the development of venous thromboembolism (VTE),<sup>129,130</sup> most of the literature supporting specific pharmacomechanical thromboprophylaxis regimens have been from mixed surgical populations.<sup>123,129</sup> Compiling data from  $>100$  RCTs of mixed abdominal surgery populations

comparing low-dose unfractionated heparin, low-molecular-weight heparin, and no chemoprophylaxis, the American College of Chest Physicians guidelines on the prevention of VTE in nonorthopedic surgical patients noted a significant reduction in VTE when prophylactic doses of either of these anticoagulant agents are used and that bleeding complications are rare.<sup>129</sup> On a case by case basis, if the consequences of bleeding are thought to be particularly severe, mechanical prophylaxis should be used until the risk abates and chemoprophylaxis can be initiated postoperatively.

If epidural or other neuraxial anesthesia is planned, it should be placed before the administration of the preoperative dose of venous thromboembolism chemoprophylaxis.<sup>104,105</sup> Chemoprophylaxis is safe throughout the hospitalization when the epidural is in place but must be held before epidural removal.

*Recommendation:* Calf-length intermittent compression devices combined with either low-dose unfractionated heparin or low-molecular-weight heparin starting immediately before surgery should be used routinely and continued at least throughout the hospitalization.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

#### 8. Preanesthetic sedative and analgesia medication.

Preoperative anxiety is frequent and can lead to increased perioperative analgesic requirements. Multiple prospective and retrospective cohort studies have demonstrated benzodiazepines can cause neurocognitive impairment and have unwanted sedative effects.<sup>131-134</sup> Therefore, sedative or anxiolytic drugs should be avoided. This is especially important for patients aged >65 years because the American Geriatric Society has made a strong recommendation, with moderate evidence, against the use of benzodiazepines for older adults.<sup>132</sup>

Preanesthetic analgesia administered orally on the day of surgery has generally been accepted as a part of a multimodal opioid-sparing perioperative analgesia strategy. The most frequently used drugs include acetaminophen, nonsteroidal anti-inflammatory drugs (NSAIDs), and gabapentinoids, all of which have been shown to decrease postoperative pain scores, facilitate opioid-sparing pain management, and can be administered cost-effectively.<sup>134,135</sup> However, although rare, the use of gabapentinoids was associated with serious respiratory depression, which was amplified when used in combination with opioid or other sedative drugs.<sup>136,137</sup> Caution should be taken in the dosing of NSAIDs and gabapentinoids in the vascular population, with adjustments determined by patient age and renal function. Consideration of NSAID-related bleeding risk is also important. However, no studies have been performed of aortic surgery to classify the risk for bleeding in the setting of low-dose NSAIDs used for a short duration in the perioperative period.<sup>138</sup> Two of the reported

enhanced recovery programs for aortic surgery included routine administration of NSAIDs but did not report adverse events.<sup>9,11</sup> Acknowledging the potential complications that can result from poor pain control or the overuse of opioids, it seems imprudent to exclude NSAIDs from perioperative use for all patients undergoing aortic surgery. No RCTs could be found in the literature of preanesthetic analgesia in open aortic surgery; however, many other surgical specialties have made strong recommendations, with low to moderate evidence, for the use of multimodal, opioid-sparing analgesia.

*Recommendation:* Sedatives should not be routinely used to reduce patient anxiety preoperatively.

*Quality of evidence:* C (low)

*Strength of recommendation:* Grade 1 (strong)

*Recommendation:* Preoperative administration of acetaminophen, NSAIDs, and gabapentinoids should be routinely used as a part of a multimodal opioid-sparing analgesia strategy.

*Quality of evidence:* C (low)

*Strength of recommendation:* Grade 1 (strong)

#### 9. Antimicrobial prophylaxis and skin preparation.

Preoperative intravenous prophylactic antibiotics should be administered immediately before (within 30-60 minutes before) open aortic aneurysm repair, with prophylactic antibiotics continued for not >24 hours postoperatively based on a meta-analysis of RCTs.<sup>56,139,140</sup> A meta-analysis of RCTs of antibiotic regimens and dosing did not demonstrate a significantly greater reduction in wound infection among comparable regimens of first- or second-generation cephalosporins, penicillin/β-lactamase inhibitors, aminoglycosides, or the glycopeptide agents vancomycin or teicoplanin.<sup>139</sup> Extrapolated from cardiac and spine surgery, the intraoperative antibiotic should be redosed within two serum half-lives of the agent<sup>141-143</sup> and can be additionally recommended for substantive procedural blood loss.<sup>144</sup>

*Recommendation:* Prophylactic intravenous antibiotic dosing should begin 30 to 60 minutes preoperatively, with redosing intraoperatively within two serum half-lives of the antimicrobial agent used or with substantive intraoperative blood loss and extending no more than 24 hours postoperatively.

*Quality of evidence:* A (high)

*Strength of recommendation:* Grade 1 (strong)

**10. Prevention of nausea and vomiting.** The prevention and treatment of postoperative nausea and vomiting (PONV) is an important component of perioperative care and one of the most common adverse events. The incidence of nausea and vomiting has been estimated to be 50% and 30% in the general surgical population, respectively, and might approach 80% for patients with

a high risk of PONV.<sup>145</sup> The risk factors for PONV include female sex, younger age, a history of PONV or motion sickness, nonsmoking, general (volatile) anesthesia, the use of postoperative opioids, and the duration of anesthesia.<sup>146</sup> The most commonly used PONV risk scoring systems are the scores by Apfel et al<sup>147</sup> and Koivuranta et al.<sup>148</sup> It is important to use PONV mitigation strategies such as minimizing the use of nitrous oxide and volatile anesthetics,<sup>149</sup> using regional anesthesia such as peripheral nerve blocks and neuraxial anesthesia,<sup>150</sup> and including opioid-sparing, multimodal analgesia techniques.<sup>151,152</sup>

Significant high-quality evidence is available from multiple RCTs supporting the use of multimodal prophylaxis and treatment of PONV that can be generalized to major vascular surgery.<sup>153</sup> Pharmacologic agent classes and other interventions available at present include 5-HT<sub>3</sub> receptor antagonists, corticosteroids, antihistamines, dopamine antagonists, propofol anesthesia, natural killer-1 receptor antagonists, anticholinergics, and acupuncture.<sup>154</sup> For rescue treatment, one should use an antiemetic from a different class than had been used for prophylaxis.

*Recommendation:* A risk assessment for PONV should be performed, multimodal PONV prophylaxis according to the findings from the assessment should be routinely used, and a PONV rescue should use a different class of antiemetic.

*Quality of evidence:* A (high)

*Strength of recommendation:* Grade 1 (strong)

## INTRAOPERATIVE RECOMMENDATIONS

### 11. Standard anesthetic protocol

Patients undergoing open aortic repair will have a high prevalence of comorbidities that predispose them to higher rates of postoperative complications and delayed recovery from surgery. The key components and aims of a standard anesthetic protocol are to use short-acting anesthetic agents and lung-protective ventilation and to achieve complete reversal of neuromuscular blockade as defined by appropriate monitoring. This allows for reversal for the patient with minimal side effects and facilitates early mobilization and feeding. A standardized multimodal anesthetic protocol should be considered in the aim to minimize the effects of anesthesia and surgery in open aortic repair and enhance recovery.

#### 11a. Neuromuscular blocking agents and monitoring.

Based on analyses of registries and other retrospective studies that have demonstrated similar results over the decades, the use of neuromuscular blocking agents should be monitored with quantitative methods (eg, acceleromyography) to ensure adequate intraoperative muscle relaxation and to prevent the delayed return of full neuromuscular function at the end of surgery. Approximately 5% of patients will experience a major pulmonary

complication after noncardiac surgery, and inadequate reversal of neuromuscular blockade increases the risk of postoperative pulmonary complications.<sup>155-158</sup> A matched cohort analysis found that selective relaxant binding agents provide a rapid and complete reversal of common steroidal neuromuscular blocking agents without adverse effects found with other drugs.<sup>158</sup> The benefits of these new agents of neuromuscular blockade reversal include fewer pulmonary complications, reduced LOS, and reduction in 30-day unplanned readmission after major abdominal surgery.<sup>159,160</sup>

*Recommendation:* Neuromuscular monitoring should be used to ensure adequate intraoperative muscle relaxation and full reversal of neuromuscular blockade before extubation.

*Quality of evidence:* A (high)

*Strength of recommendation:* Grade 1 (strong)

**11b. Cardiovascular monitoring.** Open aortic surgery has been associated with swings in hemodynamics owing to the potential for blood loss and cross-clamping and unclamping the aorta in a surgical cohort, also known to have a high prevalence of occult cardiovascular disease. Many of the changes in blood pressure and cardiovascular tone occurring during aortic surgery are well described and should be anticipated. Invasive arterial and central venous monitoring, coupled with noninvasive cardiac output monitoring, are useful in mitigating labile hemodynamics.<sup>161,162</sup> The use of transesophageal echocardiography has been well described in noncardiac surgery, in particular, its utility in the assessment of intraoperative volume status, ventricular function, and regional wall motion abnormalities.<sup>163,164</sup> These capabilities are likely to provide useful clinical information when considering the most appropriate intraoperative hemodynamic intervention at any given time; however, the specific use of transesophageal echocardiography in open aortic operations has not been studied in depth nor has any clinical outcome benefit ever been described. No evidence has shown that pulmonary artery catheters are of benefit in aortic surgery.<sup>161,162,165</sup>

Changes in mean arterial pressure, cardiac output, and heart rate associated with clamping and unclamping of the aorta make maintaining hemodynamics and intraoperative fluid therapy much more challenging in aortic cases compared with other major abdominal operations. The judicious use of vasodilators could be necessary in those with cross-clamping above the renal arteries; however, increasing the depth of anesthesia for a period after clamping will often be all that is necessary for uncomplicated cases. It is ideal to maintain the baseline blood pressure during cross-clamping to maximize collateral flow to the distal tissues. The strategies to be considered before the release of the cross-clamp include, but are not limited to, increasing the minute ventilation, reducing

the depth of anesthesia, titrating up the doses of the vasopressor or inotrope, volume loading, and sequential release of the iliac clamps.

An association exists between myocardial injury during noncardiac surgery and the intraoperative mean arterial pressure (MAP). Even relatively short periods of a low MAP have been associated with a significantly increased risk of myocardial injury.<sup>166</sup> In a prospective cohort study of vascular surgery, a 40% MAP reduction from baseline for  $\leq 30$  minutes was shown to be a significant risk.<sup>167</sup> A large retrospective cohort study (n = 57,315) reported that although relative reductions in blood pressure were a useful measure, any decrease in MAP  $< 65$  mm Hg should be avoided, because this cutoff was strongly associated with an increased rate of myocardial injury.<sup>168</sup> Maintaining MAP target levels has great importance in this patient population given the effect of aortic cross-clamping and unclamping on distal tissue and end-organ perfusion and the associated deleterious effects of ischemia–reperfusion injury during open aortic operations.

*Recommendation:* Invasive cardiovascular monitoring with an arterial line is essential. Also, the MAP should be maintained to near baseline values and at or  $> 65$  mm Hg.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

### 11c. Renal protection.

**11c-i. Ischemic preconditioning.** Much interest has ensued in the use of specific techniques to reduce organ failure or impairment after aortic surgery. One such example is ischemic preconditioning; however, this might actually cause harm.<sup>169</sup> A meta-analysis of nine trials of aortic surgery with 599 patients, of whom 336 had undergone open repair, concluded again that no benefit of renal ischemic preconditioning has been reported regarding mortality, kidney injury requiring renal support, and myocardial infarction.<sup>170</sup>

**11c-ii. Renal protective medications.** Two small RCTs of AAA patients and a meta-analysis of five small RCTs of other major cardiac and abdominal surgery have been conducted to evaluate the use of low-dose dopamine and fenoldopam on renal function in different clinical contexts. However, it appears that neither offers an advantage to euvolemic patients after elective abdominal aortic surgery.<sup>171-173</sup> The Kidney Disease: Improving Global Outcomes committee advised against the use of fenoldopam in 2012.<sup>174</sup> Similarly, RCTs have not demonstrated any clinical reductions in the incidence of renal failure with the use of mannitol.<sup>175</sup> A retrospective analysis of a single-center cohort of 169 patients who had undergone suprarenal clamping during AAA repair noted a renal protective

effect for 0.5 g/kg mannitol given before clamping.<sup>176</sup> However, a small retrospective series from the Vascular Quality Initiative registry did not note any differences in renal outcomes when stratified by mannitol administration.<sup>177</sup>

*Recommendation:* Adjunctive renal protective medications and the methods of ischemic preconditioning of any type can cause harm and should not be used for infrarenal repairs.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

**11d. Cell salvage.** The use of blood conservation strategies and, in particular, cell salvage is an established part of clinical practice and was reported in a meta-analysis of four RCTs with 292 patients to be associated with a reduced LOS in critical care, overall hospital LOS, and the use of allogeneic red blood cell transfusion, specifically for patients undergoing open AAA repair.<sup>178,179</sup> Citing these same studies, the Association of Anesthetists 2018 guidelines supported the routine use of red blood cell salvage during open AAA surgery and any surgery in adults if the blood loss could be expected to exceed 500 mL.<sup>180</sup>

*Recommendation:* Red blood cell salvage should be used.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

**11e. Lung protective ventilation.** A number of intraoperative techniques (low tidal volumes [typically 6-8 mL/kg]), the use of positive end-expiratory pressure [typically 5-7 cm H<sub>2</sub>O], and regular intraoperative recruitment maneuvers) can reduce the incidence of postoperative pulmonary complications. However, data are lacking for open aortic surgery. A recent systematic review and meta-analysis of 95 RCTs enrolling 18,062 patients reported that in enhanced recovery programs, lung-protective ventilation, goal-directed fluid therapy, and thoracic epidural analgesia are each independently associated with the reduced rates of pulmonary complications after major abdominal surgery.<sup>181</sup> Moderate quality evidence within this meta-analysis reported significant reductions in postoperative pulmonary complications for patients who had received intraoperative lung protective ventilation strategies. Owing to the high statistical heterogeneity, it is not possible to draw firm conclusions. No studies have reported evidence of harm using this strategy, and, as such, it is reasonable to adopt lung-protective ventilation strategies during open AAA surgery until further studies are undertaken and reported.<sup>182,183</sup>

*Recommendation:* Lung protective strategies should be used.

*Level of evidence:* C (low)

*Strength of recommendation:* Grade 2 (weak)

**11f. Epidural analgesia.** RCTs have compared general anesthesia with intraoperative epidural anesthesia to general anesthesia with systemic opioids with respect to reported pain, major complications, and death after open AAA surgery.<sup>184-190</sup> These studies were included in a Cochrane review in 2016, which reported no differences in 30-day mortality after open AAA surgery. However, they found low to moderate quality evidence of a reduced rate of myocardial infarction, time to extubation, LOS in critical care, and postoperative respiratory failure, favoring epidural analgesia.<sup>191</sup> The recently published National Institute for Health and Care Excellence clinical guidance on AAA management used stricter inclusion criteria and included fewer trials than those in the Cochrane group but also recommended that anesthesiologists consider using epidural analgesia in this high-risk patient group.<sup>169</sup>

One small RCT compared thoracic epidural infusion and continuous local wound infiltration in a "fast-track" abdominal aortic surgery cohort and found that the local analgesia group had required more oral and intravenous analgesics but had had a similarly short time to ambulation, regular diet, and hospital discharge.<sup>189</sup> No other data were available to make a recommendation regarding the use of transversus abdominis plane blocks or other locoregional anesthesia techniques (eg, rectus sheath and wound catheters) in open aortic surgery, although this is an area of rapidly advancing scientific study and recommendations could be possible in the future.

*Recommendation:* Mid-thoracic (T6-T9) epidural analgesia is recommended intraoperatively and should be continued postoperatively as an infusion or patient-controlled analgesia using a combination of local anesthetic and an opioid.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

## 12. Body temperature management

Perioperative hypothermia (core body temperature <36°C) results from cold operating room temperatures, patient exposure, and anesthesia-related impairment of central thermoregulatory control. Hypothermia leads to cardiac arrhythmia, coagulopathy, altered levels of consciousness, decreased drug metabolism, impaired renal function, a leftward shift in the hemoglobin oxygen saturation curve, resulting in reduced oxygen delivery, prolonged wound healing, and an increased incidence of SSI. From a combination of RCTs and prospective and retrospective cohorts across many surgical specialties, evidence has shown that maintaining normothermia reduces these undesired outcomes.<sup>192</sup> Strategies to maintain normothermia include preoperative warming (10-30 minutes before the induction of anesthesia), intraoperative warming

with a forced air heating blanket, and the use of fluid warmers and prewarmed irrigation.<sup>192</sup>

*Recommendations:* Multiple strategies should be used to maintain normothermia, including prewarming and active warming of patients intraoperatively.

*Quality of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

## 13. Drainage of surgical site strategy

Drains are not routinely used during elective transabdominal aortic surgery because these are sterile cases that do not involve the intestinal or genitourinary systems. Postoperative fluid collections will most be often hematomas, which will be ineffectively drained by small percutaneous drains. Retroperitoneal aortic surgery can result in a higher rate of lymphatic disruption. Drains have, anecdotally, been used in this approach; however, no published data were found on the benefit or harm of this practice.

In the management of groin incisions, a Cochrane review of three RCTs found no evidence demonstrating that the usage of drains will decrease groin incision complications.<sup>193</sup> The use of specific closure techniques and dressings has some increasing evidence. A single-center RCT and one cohort study of 115 patients undergoing vascular operations with a groin incision demonstrated lower rates of SSI in high-risk, but not average-risk, patients when negative pressure wound therapy was used.<sup>194,195</sup> However, a different single-center RCT of high-risk patients with groin incisions did not find a difference in SSIs when stratified by the use of negative pressure wound therapy, although the study was underpowered owing to a lower than expected rate of SSIs.<sup>196</sup>

*Recommendation:* Drains should not be routinely used in the surgical wound.

*Level of evidence:* C (low)

*Strength of recommendation:* Grade 2 (weak)

## POSTOPERATIVE RECOMMENDATIONS

**14. Multimodal analgesia and opioid reduction.** A standardized, perioperative multimodal antinociceptive protocol in enhanced recovery pathways results in adequate postoperative pain relief and improved outcomes.<sup>197-199</sup> However, the results from well-designed studies of open aortic surgery have been inconclusive or are lacking.

Acetaminophen and NSAIDs, including selective COX-2 (cyclooxygenase-2) inhibitors, are a basic part of postoperative multimodal pain management and are used widely, either orally or intravenously.<sup>198,200,201</sup> Acetaminophen is an antipyretic, and its analgesic activity is additive to NSAIDs and opioids. NSAIDs should be part of a

multimodal strategy after open aortic surgery unless the patient has specific contraindications for its use. Several RCTs of other surgical specialties have also investigated other nonopioid postoperative analgesia, including gabapentin,  $\alpha_2$ -agonists, S-ketamine, magnesium sulfate,  $\beta$ -blockade, and high dose steroids, which all resulted in a reduction in pain.<sup>202-208</sup>

Opioids have remained the most frequently prescribed class of drugs globally.<sup>198</sup> However, it has also been recognized that patients are at a high risk of opioid misuse and addiction in the long term.<sup>209,210</sup> In the short term, opioids have also been related to many undesirable side effects that slow recovery, specifically including delirium, which is of particular relevance to the aortic surgery population.<sup>211,212</sup> Oral or intravenous opioids might be less effective in treating acute postoperative pain following open aortic surgery compared with epidural analgesia.<sup>213</sup> However, in a single-center RCT of 168 patients undergoing open aortic surgery, patients with patient-controlled epidural analgesia had had similar outcomes and time to recovery compared with the patients with patient-controlled intravenous narcotic analgesia.<sup>190</sup>

*Recommendation:* Multimodal analgesic regimens should be routinely used to improve pain control and minimize opioid consumption.

*Quality of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

**15. Nasogastric drainage.** Meta-analyses of nonrandomized studies of patients undergoing gastrectomy or other abdominal surgery have demonstrated that the routine use of nasogastric tubes is associated with delayed oral intake and return of bowel function.<sup>214-216</sup> For operations that also involve foregut mobilization, the 2019 ERAS guidelines for pancreatoduodenectomy recommend against the routine use of nasogastric tubes and advocate removing them before the end of the anesthesia.<sup>90</sup> It is reasonable to generalize this practice of removing the nasogastric tube at the time of extubation to aortic patients who undergo duodenal mobilization, but not resection. The routine use of nasogastric tubes has been shown in large Cochrane analysis of RCTs to cause pneumonia.<sup>217</sup>

*Recommendation:* The routine use of postoperative nasogastric tubes should be avoided.

*Level of evidence:* A (high)

*Strength of recommendation:* Grade 1 (strong)

**16. Oral feeding.** Traditional signs of bowel function such as the presence of bowel sounds and passage of flatus are unreliable indicators of a return of gastric and intestinal function. In a cohort study of patients undergoing elective, open AAA repair, the median time to normal gastric emptying was  $18 \pm 7$  hours, indicating that little

reason exists to withhold solid food from patients after 24 hours.<sup>218</sup> Early enteral nutrition has been shown to reduce infectious complications and to be protective against the development of intestinal ischemia and stress ulcers in a prospective cohort study of AAA patients in critical care units.<sup>219</sup> Postoperatively, patients should be encouraged to reestablish oral nutrition as early as possible, allowing patients to eat what they are comfortable with and consume oral nutritional supplements daily. Oral intake of solid foods has been shown to be possible within 24 hours in multiple "fast-track" observational studies of patients recovering from open aortic surgery in Europe.<sup>8,11,12,220,221</sup>

*Recommendation:* An early return to a normal diet and oral nutritional supplements should be promoted.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

**17. Fluid therapy.** Fluid management has been a central tool of postoperative therapy in aortic surgery, and appropriate fluid replacement is needed to maintain a euvolemic state and avoid postoperative complications, including acute kidney injury,<sup>222,223</sup> or the sequelae of fluid overload, including heart failure, decreased oxygenation, and prolonged ileus.<sup>224-226</sup> Ideally, patients will leave the operating room in a euvolemic state, rendering fluid boluses, on the one hand, or diuretics, on the other hand, unnecessary. One's best clinical judgment must be used to correct the fluid balance in patients who leave the operating room under- or over-resuscitated. However, one must remember that significant variability exists in the subjective fluid balance assessment between providers and within the same provider, which is strong predictor of fluid overload.<sup>227,228</sup>

If euvolemic patients are able to drink fluids, the use of intravenous fluids might not be needed to maintain euvolemia.<sup>8</sup> However, if needed, restrictive fluid therapy has been more beneficial than liberal fluid replacement. Three RCTs with a total of 125 aortic surgery patients without existing severe congestive heart failure or chronic kidney disease demonstrated that the group given  $\leq 1.5$  L/d had significantly fewer complications, a lower risk of cardiopulmonary complications, and shorter ICU and total hospital LOSs than the liberal fluid therapy group.<sup>229-231</sup> A Cochrane review reported no evidence for the use of a specific fluid or evidence to support colloids vs crystalloids for patients undergoing abdominal aortic surgery.<sup>232</sup>

*Recommendation:* If intravenous fluids are indicated, postoperative fluid therapy should be restricted to 1.5 L/d.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

**18. Urinary drainage.** Accurate measurement of urine output is necessary in the immediate postoperative

period, because acute kidney injury has been estimated to develop in  $\leq 15\%$  of patients after infrarenal clamping and 21% of patients after suprarenal or supraceliac clamping.<sup>177,233-235</sup> Because of concerns of catheter-associated urinary tract infection and guidance from the U.S. Centers for Disease Control and Prevention, the use of prolonged urinary catheters has decreased.<sup>236</sup> As such, urinary catheters placed for surgical and postoperative monitoring should be removed as soon as possible, preferably within 24 hours.<sup>236</sup> The presence of a thoracic epidural should not preclude removal of urinary drainage catheters. However,  $\sim 10\%$  of patients overall and 25% of patients with benign prostatic hyperplasia will require catheter replacement because of urinary retention.<sup>237,238</sup>

*Recommendation:* The early removal of urinary drainage catheters is recommended.

*Level of evidence:* C (low)

*Strength of recommendation:* Grade 1 (strong)

**19. Glycemic control.** Poor glycemic control is an independent risk factor for adverse events in vascular patients with or without a diagnosis of diabetes.<sup>239,240</sup> In a cohort study of  $>1000$  patients undergoing vascular surgery, postoperative hyperglycemia was strongly associated with 11 different negative outcomes, including 30-day mortality, myocardial infarction, acute renal failure, stroke, wound complications, and a return to the operating room.<sup>240</sup> Nondiabetic patients with hyperglycemia demonstrated an even stronger association with these negative surgical outcomes.<sup>240</sup>

In a multicenter RCT of 880 nondiabetic patients undergoing major abdominal surgery, preoperative carbohydrate loading prevented postoperative hyperglycemia and significantly reduced the number of patients requiring insulin to maintain a glucose level  $<180$  mg/dL.<sup>110</sup> A single-center RCT of 236 vascular patients undergoing open AAA repair, lower extremity bypass surgery, or limb amputation compared continuous insulin infusion to maintain glucose levels at 100 to 150 mg/dL to intermittent insulin boluses to maintain glucose levels  $>150$  mg/dL during the first 48 hours after surgery. Insulin infusion was protective against a composite end point of all-cause death, myocardial infarction, and congestive heart failure, and those patients did not experience a statistically significant different rate of hypoglycemia.<sup>239</sup> Careful postoperative glycemic control with an emphasis on limiting extreme variability in glucose levels decreased the incidence of adverse cardiac events.<sup>239</sup> However, caution is advised because overly aggressive glucose control causing hypoglycemia can increase mortality.<sup>241,242</sup> No RCT data for a serum glucose threshold for initiating intraoperative insulin treatment, and no perioperative glucose targets have been agreed on for patients undergoing aortic operations.<sup>243</sup>

*Recommendation:* Glycemic control should be maintained, with intention of avoiding hypoglycemia.

*Level of evidence:* B (moderate)

*Strength of recommendation:* Grade 1 (strong)

**20. Early mobilization.** No high-quality studies regarding early mobility and open aortic surgery; however, many studies of ERAS-like pathways for open aortic surgery effectively used early mobility as a part of their care bundles.<sup>8</sup> Extrapolating from other specialties, early mobility after aortic surgery likely decreases complications and improves patient outcomes.<sup>244,245</sup> Successful mobilization strategies require a multidisciplinary approach, objective evaluation of patient safety and an ability for early ambulation, and tracking progress with a formal mobility scale.<sup>246</sup> An early mobility protocol with at least twice-daily assessments increases mobility and decreases delirium and LOS.<sup>246</sup> Those who are more physically fit preoperatively will be able to achieve postoperative mobility goals more quickly<sup>95</sup>; however, all patients should have a tailored, safe plan for ambulation.<sup>246</sup> If patients are not progressing, the causes of failure should be analyzed by the multidisciplinary team, including physical therapists, and corrective measures should be instituted quickly.

*Recommendation:* The use of a formal plan for early mobilization should be instituted, with early physical therapy involvement, when needed.

*Level of evidence:* C (low)

*Strength of recommendation:* Grade 1 (strong)

**21. Discharge education.** It is reasonable to believe that the standard discharge criteria for many general surgery operations are applicable to aortic surgery patients, including tolerance of oral intake, recovery of lower gastrointestinal function, adequate pain control, ability to mobilize and perform self-care, and no evidence of complications or untreated medical problems.<sup>247</sup> Up to 30% of aortic patients might have delirium or transient advanced mental impairment that complicates or delays discharge readiness.<sup>248</sup>

Patients undergoing vascular surgery have had 30-day readmission rates as high as 20% and up to one half of readmissions will be deemed avoidable.<sup>249-251</sup> In line with fundamental ERAS philosophy, a published conceptual model of preventing readmissions in vascular surgery that includes patient education and timely communication from the preoperative phase to the postoperative phase of care.<sup>252</sup> A Cochrane review of two RCTs concluded that written and verbal information on specific care instructions, medication lists, and when to seek medical attention at discharge significantly increased patient knowledge and satisfaction.<sup>253,254</sup> The data on follow-up postoperative telephone calls has

been mixed. A Cochrane review of 33 studies of mixed methods from different specialties was inconclusive; however, no adverse effects of the invention were reported.<sup>255</sup> In-person visits might be more beneficial. Among patients experiencing a complication after open thoracic aneurysm repair, having a visit with a primary care physician within 30 days of the operation reduced the readmission rates from 35% to 20%.<sup>256</sup>

**Recommendation:** Both written and verbal discharge education should be used, with follow-up telephone calls within 24 to 48 hours after hospital discharge, and clinic follow-up visits should be scheduled with both the surgeon and the primary care physician at least once by 30 days postoperatively.

**Level of evidence:** C (low)

**Strength of recommendation:** grade 1 (strong)

**22. Audit of outcomes.** Previous implementation of ERAS protocols in other surgical disciplines has led to a reduction in complications, shorter LOSs, and improved cost-savings. The analysis of the literature on ERAS audits has been based almost exclusively on systematic reviews, and prospective studies on this topic have yet to be developed. However, one prospective analysis comparing self-declared ERAS with non-ERAS hospitals demonstrated that having an ERAS protocol in place is not enough to improve patient outcomes.<sup>257</sup> Other surgical specialties have demonstrated an association between increasing ERAS compliance and decreased surgical complications and LOS.<sup>258,259</sup>

Interactive auditing and reporting will identify areas of noncompliance, improve tracking of outcomes, and facilitate continuous quality improvement related to specific patient populations, care settings, and/or logistical issues.<sup>258</sup> An observational study of >500 gynecologic patients found that using a compliance audit tool increased compliance from 56% to 77% and was associated with significant reductions in LOS.<sup>260</sup> Another single-center pre–post study of gynecologic patients demonstrated that structured ERAS implementation, including interactive auditing, resulted in better ERAS compliance and a shorter time to functional recovery and hospital discharge.<sup>261</sup> Similar results were found at 13 centers that had participated in an international ERAS database for colorectal surgery; however, interestingly, 2 years after initiating the ERAS programs, auditing revealed that implementation compliance had decreased, which correlated with an increase in the LOS and complication rates.<sup>262</sup> To improve the quality of ERAS auditing and reporting, the ERAS USA and ERAS Society have published the Reporting on ERAS Compliance, Outcomes, and Elements Research Checklist.<sup>7</sup> This tool delineates the best practices for reporting clinical pathways and describing compliance. Clinicians are encouraged to use auditing tools; examples include the ERAS Interactive Audit System (available at: <http://erassociety.org/interactive-audit/>) and the Agency for

Healthcare Research and Quality safety program for improving surgical care and recovery (available at: <https://www.ahrq.gov/hai/tools/enhanced-recovery/index.html>).

**Recommendation:** The use of routine auditing of ERAS care processes and feedback to all staff involved is necessary.

**Level of evidence:** C (low)

**Strength of recommendation:** Grade 1 (strong)

## CARE ELEMENTS WITH CONFLICTING OR INSUFFICIENT DATA FOR A RECOMMENDATION

Some traditional clinical practices are in use for which the literature does not allow firm recommendations; therefore, they were not included in our consensus statement. Nevertheless, because of their common use in many institutions, the writing group has summarized the current state of knowledge.

**Preoperative medical risk assessment.** A number of commonly used practices are of limited or no benefit in routine medical risk assessment. Preoperative routine chest radiography cannot be recommended because it has no supporting scientific evidence. It can be selectively used for patients with new or unstable cardiopulmonary signs or symptoms and for patients at increased risk of postoperative pulmonary complications if the results will alter management (ie, postpone surgery).<sup>263-265</sup> In a large, prospective cohort study, the results from preoperative pulmonary function studies did not correlate with the postoperative outcomes and cannot be recommended as a routine part of a preoperative workup.<sup>266</sup>

Sources of infection, apart from skin and the operative environment, should be considered, such as elimination of potential septic dental sources  $\geq 2$  weeks before implantation of an aortic prosthesis.<sup>56</sup> The predictive value of routine urinalysis in asymptomatic patients is poor<sup>267,268</sup> but must be weighed against the potential infectious risk to prosthetic material.<sup>56</sup> Although these measures make sense, no solid scientific background is available for a firm and specific recommendation.

Hematologic disorders, specifically thrombocytopenia, should be evaluated preoperatively, preferably by a hematologist, because of the associated bleeding and mortality risk.<sup>269,270</sup>

**Skin preparation.** Based on meta-analysis of a small number of RCTs with heterogeneous surgical procedures and quasi-experimental cohort studies of vascular operations, hair clipping should be performed immediately preoperatively.<sup>271-273</sup> Preoperative skin preparation should be standardized, with the operative area being scrubbed the day before and immediately before surgery with a chlorhexidine solution. This was based on RCTs of heterogeneous clean-contaminated surgeries, which indicated a benefit. However, conflicting evidence was

found in smaller RCTs and prospective observational studies of vascular reconstructions that did not demonstrate benefit.<sup>274-276</sup>

**Intraoperative fluid therapy.** No reliable data are available to guide fluid management during open aortic surgery, and the need for research is obvious. The use of goal-directed fluid therapy has been recommended for many other major abdominal surgeries. However, fluid management during open aortic surgery is particularly complicated because of cardiac disease and cross-clamping, which causes extreme disruptions in cardiac stress and fluid distribution in the body.

## SUMMARY AND CONCLUSIONS

This consensus statement represents the most recent evidenced-based recommendations from the ERAS Society guideline group and the SVS for the perioperative management of patients undergoing open aortic vascular surgery.

These first ERAS recommendations for elective open aortic surgery are important in summarizing the large volume of heterogeneous studies across all ERAS elements for operations performed for either aortic aneurysm disease or aortoiliac occlusive disease. The authors' recommendations provided in this intersocietal consensus statement are in accordance with the methods set out by the ERAS Society and SVS and are based on the synthesis of an objective assessment of the best available evidence in open aortic surgery, other surgical disciplines, and consensus of the guideline development group. The available evidence was graded and potential harm from the use of the care element was also considered in the recommendations; therefore, strong recommendations could have been reached from low-quality or conflicting data and vice versa. Likewise, this method clearly accounts for, and indicates where recommendations have been made, based on data extrapolated from other surgical specialties.

The present consensus statement highlights the numerous research gaps that exist. Although the results from the few clinical studies available of cohorts of patients with aortic disease are promising, studies of high methodologic quality are needed. The lines of research to be developed could include any of the ERAS elements or combined elements but are likely to be particularly needed for preadmission optimization, prehabilitation, preoperative carbohydrate loading for those with diabetes, preanesthetic sedatives and analgesia, conditions for postoperative urinary catheter removal, early mobility protocols, postdischarge education, and auditing practices.

Our main purpose was to define the current standards based on the available literature and to enable multidisciplinary teams to implement these procedures in their practice to investigate to what extent they can improve

outcomes.<sup>1,277</sup> As demonstrated in other specialties, using similar evidence-based guidelines and consensus statements to form care teams to enhance patient optimization and shared decision-making, focus on physiologic stress minimization, and reduce variations in postoperative convalescence promises to improve the safety and outcomes for patients undergoing open aortic operations.

## REFERENCES

1. Ljungqvist O, Scott M, Fearon KC. Enhanced recovery after surgery: a review. *JAMA Surg* 2017;152:292-8.
2. Kehlet H. Multimodal approach to control postoperative pathophysiology and rehabilitation. *Br J Anaesth* 1997;78:606-17.
3. Fearon KC, Ljungqvist O, Von Meyenfeldt M, Revhaug A, Dejong CH, Lassen K, et al. Enhanced recovery after surgery: a consensus review of clinical care for patients undergoing colonic resection. *Clin Nutr* 2005;24:466-77.
4. Brindle M, Nelson G, Lobo DN, Ljungqvist O, Gustafsson UO. Recommendations from the ERAS Society for standards for the development of enhanced recovery after surgery guidelines. *BJS Open* 2020;4:157-63.
5. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924-6.
6. Murad MH, Montori VM, Sidawy AN, Ascher E, Meissner MH, Chaikof EL, et al. Guideline methodology of the Society for Vascular Surgery including the experience with the GRADE framework. *J Vasc Surg* 2011;53:1375-80.
7. Elias KM, Stone AB, McGinigle K, Tankou JI, Scott MJ, Fawcett WJ, et al. The Reporting on ERAS Compliance, Outcomes, and Elements Research (RECOVER) Checklist: a joint statement by the ERAS((R)) and ERAS((R)) USA Societies. *World J Surg* 2019;43:1-8.
8. McGinigle KL, Eldrup-Jorgensen J, McCall R, Freeman NL, Pascarella L, Farber MA, et al. A systematic review of enhanced recovery after surgery for vascular operations. *J Vasc Surg* 2019;70:629-40.e1.
9. Tatsuiishi W, Kohri T, Koderia K, Asano R, Kataoka G, Kubota S, et al. Usefulness of an enhanced recovery after surgery protocol for perioperative management following open repair of an abdominal aortic aneurysm. *Surg Today* 2012;42:1195-200.
10. Feo CV, Portinari M, Tsolaki E, Romagnoni G, Verri M, Camerani S, et al. The effect of an enhanced recovery program in elective retroperitoneal abdominal aortic aneurysm repair. *J Vasc Surg* 2016;63:888-94.
11. Muehling B, Schelzig H, Steffen P, Meierhenrich R, Sunder-Plassmann L, Orend KH. A prospective randomized trial comparing traditional and fast-track patient care in elective open infrarenal aneurysm repair. *World J Surg* 2009;33:577-85.
12. Brustia P, Renghi A, Aronici M, Gramaglia L, Porta C, Musiani A, et al. Fast-track in abdominal aortic surgery: experience in over 1,000 patients. *Ann Vasc Surg* 2015;29:1151-9.
13. Murphy MA, Richards T, Atkinson C, Perkins J, Hands LJ. Fast track open aortic surgery: reduced postoperative stay with a goal directed pathway. *Eur J Vasc Endovasc Surg* 2007;34:274-8.
14. Forsmo HM, Erichsen C, Rasdal A, Tvinnereim JM, Körner H, Pfeffer F. Randomized controlled trial of extended perioperative counseling in enhanced recovery after colorectal surgery. *Dis Colon Rectum* 2018;61:724-32.
15. de Aguiar-Nascimento JE, Leal FS, Dantas DC, Anabuki NT, de Souza AM, Silva ELVP, et al. Preoperative education in cholecystectomy in the context of a multimodal protocol of perioperative care: a randomized, controlled trial. *World J Surg* 2014;38:357-62.
16. Gräwe JS, Mirow L, Bouchard R, Lindig M, Hüppe M. [Impact of preoperative patient education on postoperative pain in consideration of the individual coping style]. *Schmerz (Berlin, Germany)* 2010;24:575-86.
17. Muehling BM, Ortlieb L, Oberhuber A, Orend KH. Fast track management reduces the systemic inflammatory response and organ failure following elective infrarenal aortic aneurysm repair. *Interact Cardiovasc Thorac Surg* 2011;12:784-8.

18. Baron DM, Hochrieser H, Posch M, Metnitz B, Rhodes A, Moreno RP, et al. Preoperative anaemia is associated with poor clinical outcome in non-cardiac surgery patients. *Br J Anaesth* 2014;113:416-23.
19. Musallam KM, Tamim HM, Richards T, Spahn DR, Rosendaal FR, Habbal A, et al. Preoperative anaemia and postoperative outcomes in non-cardiac surgery: a retrospective cohort study. *Lancet* 2011;378:1396-407.
20. Spahn DR, Goodnough LT. Alternatives to blood transfusion. *Lancet* 2013;381:1855-65.
21. Isbister JP, Shander A, Spahn DR, Erhard J, Farmer SL, Hofmann A. Adverse blood transfusion outcomes: establishing causation. *Transfus Med Rev* 2011;25:89-101.
22. Spahn DR, Schoenrath F, Spahn GH, Seifert B, Stein P, Theusinger OM, et al. Effect of ultra-short-term treatment of patients with iron deficiency or anaemia undergoing cardiac surgery: a prospective randomised trial. *Lancet* 2019;393:2201-12.
23. Guinn NR, Guercio JR, Hopkins TJ, Grimsley A, Kurian DJ, Jimenez MI, et al. How do we develop and implement a preoperative anemia clinic designed to improve perioperative outcomes and reduce cost? *Transfusion* 2016;56:297-303.
24. Enko D, Wallner F, von-Goedecke A, Hirschmugl C, Auersperg V, Halwachs-Baumann G. The impact of an algorithm-guided management of preoperative anemia in perioperative hemoglobin level and transfusion of major orthopedic surgery patients. *Anemia* 2013;2013:641876.
25. Goodnough LT, Maniatis A, Earnshaw P, Benoni G, Beris P, Bisbe E, et al. Detection, evaluation, and management of preoperative anaemia in the elective orthopaedic surgical patient: NATA guidelines. *Br J Anaesth* 2011;106:13-22.
26. Kozek-Langenecker SA, Ahmed AB, Afshari A, Albaladejo P, Aldecoa C, Barauskas G, et al. Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology: first update 2016. *Eur J Anaesthesiol* 2017;34:332-95.
27. Janssen TL, Steyerberg EW, van Gammeren AJ, Ho GH, Gobardhan PD, van der Laan L. Intravenous iron in a prehabilitation program for older surgical patients: prospective cohort study. *J Surg Res* 2021;257:32-41.
28. Gustafsson UO, Scott MJ, Hubner M, Nygren J, Demartines N, Francis N, et al. Guidelines for perioperative care in elective colorectal surgery: Enhanced Recovery After Surgery (ERAS) Society recommendations: 2018. *World J Surg* 2019;43:659-95.
29. Ng O, Keeler BD, Mishra A, Simpson JA, Neal K, Al-Hassi HO, et al. Iron therapy for preoperative anaemia. *Cochrane Database Syst Rev* 2019;12:CD011588.
30. Richards T, Baikady RR, Clevenger B, Butcher A, Abeysiri S, Chau M, et al. Preoperative intravenous iron to treat anaemia before major abdominal surgery (PREVENTT): a randomised, double-blind, controlled trial. *Lancet* 2020;396:1353-61.
31. U.S. Food and Drug Administration. Epogen (epoetin alfa) injection, for intravenous or subcutaneous use. Drug label. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2017/103234s5363s53661bl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/103234s5363s53661bl.pdf). Accessed September 28, 2020.
32. Stoppe C, Goetzenich A, Whitman G, Ohkuma R, Brown T, Hatzakorzian R, et al. Role of nutrition support in adult cardiac surgery: a consensus statement from an International Multidisciplinary Expert Group on Nutrition in Cardiac Surgery. *Crit Care* 2017;21:131.
33. Lomivorotov VV, Efremov SM, Boboshko VA, Nikolaev DA, Vedernikov PE, Lomivorotov VN, et al. Evaluation of nutritional screening tools for patients scheduled for cardiac surgery. *Nutrition* 2013;29:436-42.
34. Lomivorotov VV, Efremov SM, Boboshko VA, Nikolaev DA, Vedernikov PE, Deryagin MN, et al. Prognostic value of nutritional screening tools for patients scheduled for cardiac surgery. *Interact Cardiovasc Thorac Surg* 2013;16:612-8.
35. Kodama A, Takahashi N, Sugimoto M, Niimi K, Banno H, Komori K. Associations of nutritional status and muscle size with mortality after open aortic aneurysm repair. *J Vasc Surg* 2019;70:1585-93.
36. Weimann A, Braga M, Carli F, Higashiguchi T, Hubner M, Klek S, et al. ESPEN guideline: clinical nutrition in surgery. *Clin Nutr* 2017;36:623-50.
37. Lobo DN, Gianotti L, Adiamah A, Barazzoni R, Deutz NEP, Dhatriya K, et al. Perioperative nutrition: recommendations from the ESPEN expert group. *Clin Nutr* 2020;39:3211-27.
38. Moya P, Soriano-Irigaray L, Ramirez JM, Garcea A, Blasco O, Blanco FJ, et al. Perioperative standard oral nutrition supplements versus immunonutrition in patients undergoing colorectal resection in an enhanced recovery (ERAS) protocol: a multicenter randomized clinical trial (SONVI study). *Medicine* 2016;95:e3704.
39. Buzquurz F, Bojesen RD, Grube C, Madsen MT, Gogenur I. Impact of oral preoperative and perioperative immunonutrition on postoperative infection and mortality in patients undergoing cancer surgery: systematic review and meta-analysis with trial sequential analysis. *BJS Open* 2020;4:764-75.
40. Moya P, Miranda E, Soriano-Irigaray L, Arroyo A, Aguilar M-D-M, Bellon M, et al. Perioperative immunonutrition in normo-nourished patients undergoing laparoscopic colorectal resection. *Surg Endosc* 2016;30:4946-53.
41. Waitzberg DL, Saito H, Plank LD, Jamieson GG, Jagannath P, Hwang T-L, et al. Postsurgical infections are reduced with specialized nutrition support. *World J Surg* 2006;30:1592-604.
42. Jie B, Jiang Z-M, Nolan MT, Zhu S-N, Yu K, Kondrup J. Impact of preoperative nutritional support on clinical outcome in abdominal surgical patients at nutritional risk. *Nutrition* 2012;28:1022-7.
43. McClave SA, Kozar R, Martindale RG, Heyland DK, Braga M, Carli F, et al. Summary points and consensus recommendations from the North American Surgical Nutrition Summit. *JPEN J Parenter Enteral Nutr* 2013;37(Suppl):99S-105S.
44. Arya S, Kim SI, Duwayri Y, Brewster LP, Veeraswamy R, Salam A, et al. Frailty increases the risk of 30-day mortality, morbidity, and failure to rescue after elective abdominal aortic aneurysm repair independent of age and comorbidities. *J Vasc Surg* 2015;61:324-31.
45. Al Shakarchi J, Fairhead J, Rajagopalan S, Pherwani A, Jaipersad A. Impact of frailty on outcomes in patients undergoing open abdominal aortic aneurysm repair. *Ann Vasc Surg* 2020;67:100-4.
46. Arya S, Long CA, Brahmabhatt R, Shafiq S, Brewster LP, Veeraswamy R, et al. Preoperative frailty increases risk of nonhome discharge after elective vascular surgery in home-dwelling patients. *Ann Vasc Surg* 2016;35:19-29.
47. Donald GW, Ghaffarian AA, Isaac F, Kraiss LW, Griffin CL, Smith BK, et al. Preoperative frailty assessment predicts loss of independence after vascular surgery. *J Vasc Surg* 2018;68:1382-9.
48. George EL, Chen R, Trickey AW, Brooke BS, Kraiss L, Mell MW, et al. Variation in center-level frailty burden and the impact of frailty on long-term survival in patients undergoing elective repair for abdominal aortic aneurysms. *J Vasc Surg* 2020;71:46-55.e4.
49. George EL, Kashikar A, Rothenberg KA, Barreto NB, Chen R, Trickey AW, et al. Comparison of surgeon assessment to frailty measurement in abdominal aortic aneurysm repair. *J Surg Res* 2020;248:38-44.
50. Alvarez-Nebreda ML, Bentov N, Urman RD, Setia S, Huang JC, Pfeifer K, et al. Recommendations for preoperative management of frailty from the Society for Perioperative Assessment and Quality Improvement (SPAQI). *J Clin Anesth* 2018;47:33-42.
51. Partridge JS, Harari D, Martin FC, Peacock JL, Bell R, Mohammed A, et al. Randomized clinical trial of comprehensive geriatric assessment and optimization in vascular surgery. *Br J Surg* 2017;104:679-87.
52. Raats JW, van Eijnsden WA, Crolla RM, Steyerberg EW, van der Laan L. Risk factors and outcomes for postoperative delirium after major surgery in elderly patients. *PLoS One* 2015;10:e0136071.
53. Janssen TL, Steyerberg EW, Faes MC, Wijsman JH, Gobardhan PD, Ho GH, et al. Risk factors for postoperative delirium after elective major abdominal surgery in elderly patients: a cohort study. *Int J Surg* 2019;71:29-35.
54. Janssen TL, Steyerberg EW, Langenberg JCM, de Lepper C, Wielders D, Seerden TCJ, et al. Multimodal prehabilitation to reduce the incidence of delirium and other adverse events in elderly patients undergoing elective major abdominal surgery: an uncontrolled before-and-after study. *PLoS One* 2019;14:e0218152.
55. Wallbridge HR, Benoit AG, Staley D, Ediger JP, Campbell BI. Risk factors for postoperative cognitive and functional difficulties in abdominal aortic aneurysm patients: a three month follow-up. *Int J Geriatr Psychiatry* 2011;26:818-24.
56. Chaikof EL, Dalman RL, Eskandari MK, Jackson BM, Lee WA, Mansour MA, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *J Vasc Surg* 2018:2-77.

57. Wong J, Lam DP, Abrishami A, Chan MT, Chung F. Short-term pre-operative smoking cessation and postoperative complications: a systematic review and meta-analysis. *Can J Anesth* 2012;59:268-79.
58. Tonnesen H, Nielsen PR, Lauritzen JB, Moller AM. Smoking and alcohol intervention before surgery: evidence for best practice. *Br J Anaesth* 2009;102:297-306.
59. Sorensen LT. Wound healing and infection in surgery: the clinical impact of smoking and smoking cessation: a systematic review and meta-analysis. *Arch Surg* 2012;147:373-83.
60. Hughes JR, Stead LF, Hartmann-Boyce J, Cahill K, Lancaster T. Antidepressants for smoking cessation. *Cochrane Database Syst Rev* 2014;1:CD000031.
61. Oppedal K, Møller AM, Pedersen B, Tønnesen H. Preoperative alcohol cessation prior to elective surgery. *Cochrane Database Syst Rev* 2012;7:CD008343.
62. Shabanzadeh DM, Sorensen LT. Alcohol consumption increases post-operative infection but not mortality: a systematic review and meta-analysis. *Surg Infect (Larchmt)* 2015;16:657-68.
63. Bahia SS, Holt PJ, Jackson D, Patterson BO, Hinchliffe RJ, Thompson MM, et al. Systematic review and meta-analysis of long-term survival after elective infrarenal abdominal aortic aneurysm repair 1969-2011: 5 year survival remains poor despite advances in medical care and treatment strategies. *Eur J Vasc Endovasc Surg* 2015;50:320-30.
64. Fleisher LA, Fleischmann KE, Auerbach AD, Barnason SA, Beckman JA, Bozkurt B, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *Circulation* 2014;130:e278-333.
65. Lee TH, Marcantonio ER, Mangione CM, Thomas EJ, Polanczyk CA, Cook EF, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation* 1999;100:1043-9.
66. Gupta PK, Gupta H, Sundaram A, Kaushik M, Fang X, Miller WJ, et al. Development and validation of a risk calculator for prediction of cardiac risk after surgery. *Circulation* 2011;124:381-7.
67. Bertges DJ, Neal D, Schanzer A, Scali ST, Goodney PP, Eldrup-Jorgensen J, et al. The Vascular Quality Initiative cardiac risk index for prediction of myocardial infarction after vascular surgery. *J Vasc Surg* 2016;64:1411-21.e4.
68. Eslami MH, Rybin DV, Doros G, Siracuse JJ, Farber A. External validation of Vascular Study Group of New England risk predictive model of mortality after elective abdominal aorta aneurysm repair in the Vascular Quality Initiative and comparison against established models. *J Vasc Surg* 2018;67:143-50.
69. National Institute for Health and Care Excellence. Surveillance report 2018. Cardiovascular disease: risk assessment and reduction, including lipid modification. NICE guideline CG181. Available at: <https://www.nice.org.uk/guidance/cg181> 2014. Accessed December 10, 2020.
70. Pearson TA, Blair SN, Daniels SR, Eckel RH, Fair JM, Fortmann SP, et al. AHA guidelines for primary prevention of cardiovascular disease and stroke: 2002 update: consensus panel guide to comprehensive risk reduction for adult patients without coronary or other atherosclerotic vascular diseases. American Heart Association Science Advisory and Coordinating Committee. *Circulation* 2002;106:388-91.
71. Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, et al. 2016 European guidelines on cardiovascular disease prevention in clinical practice. *Rev Esp Cardiol (Engl Ed)* 2016;69:939.
72. Alagona P Jr, Ahmad TA. Cardiovascular disease risk assessment and prevention: current guidelines and limitations. *Med Clin North Am* 2015;99:711-31.
73. Bourl S, Shun-Shin MJ, Cole GD, Mayet J, Francis DP. Meta-analysis of secure randomised controlled trials of beta-blockade to prevent perioperative death in non-cardiac surgery. *Heart* 2014;100:456-64.
74. American Diabetes Association. 10. Cardiovascular disease and risk management: standards of medical care in diabetes—2019. *Diabetes Care* 2019;42(Suppl 1):S103-23.
75. Newman JD, Wilcox T, Smilowitz NR, Berger JS. Influence of diabetes on trends in perioperative cardiovascular events. *Diabetes Care* 2018;41:1268-74.
76. Martin-Gonzalez T, Pincon C, Hertault A, Maurel B, Labbe D, Spear R, et al. Renal outcomes analysis after endovascular and open aortic aneurysm repair. *J Vasc Surg* 2015;62:569-77.
77. Nguyen BN, Neville RF, Rahbar R, Amdur R, Sidawy AN. Comparison of outcomes for open abdominal aortic aneurysm repair and endovascular repair in patients with chronic renal insufficiency. *Ann Surg* 2013;258:394-9.
78. Kellum JA, Lameire N; KDIGO AKI Guideline Work Group. Diagnosis, evaluation, and management of acute kidney injury: a KDIGO summary (part 1). *Crit Care* 2013;17:204.
79. Bown MJ, Norwood MG, Sayers RD. The management of abdominal aortic aneurysms in patients with concurrent renal impairment. *Eur J Vasc Endovasc Surg* 2005;30:1-11.
80. Pouwels S, Willigendael EM, van Sambeek MR, Nienhuijs SW, Cuypers PW, Teijink JA. Beneficial effects of pre-operative exercise therapy in patients with an abdominal aortic aneurysm: a systematic review. *Eur J Vasc Endovasc Surg* 2015;49:66-76.
81. Gillis C, Gramlich L, Culos-Reed SN, Sajobi TT, Fiest KM, Carli F, et al. Third-variable effects: tools to understand who, when, why, and how patients benefit from surgical prehabilitation. *J Surg Res* 2021;258:443-52.
82. Carli F, Awasthi R, Gillis C, Baldini G, Bessissow A, Liberman AS, et al. Integrating prehabilitation in the preoperative clinic: a paradigm shift in perioperative care. *Anesth Analg* 2021;132:1494-500.
83. Arthur HM, Daniels C, McKelvie R, Hirsh J, Rush B. Effect of a pre-operative intervention on preoperative and postoperative outcomes in low-risk patients awaiting elective coronary artery bypass graft surgery: a randomized, controlled trial. *Ann Intern Med* 2000;133:253-62.
84. Sawatzky JA, Kehler DS, Ready AE, Lerner N, Boreskie S, Lamont D, et al. Prehabilitation program for elective coronary artery bypass graft surgery patients: a pilot randomized controlled study. *Clin Rehabil* 2014;28:648-57.
85. Stammers AN, Kehler DS, Afilalo J, Avery LJ, Bagshaw SM, Crocetti HP, et al. Protocol for the PREHAB study—pre-operative rehabilitation for reduction of hospitalization after coronary bypass and valvular surgery: a randomised controlled trial. *BMJ Open* 2015;5:e007250.
86. Barberan-Garcia A, Ubre M, Roca J, Lacy AM, Burgos F, Risco R, et al. Personalised prehabilitation in high-risk patients undergoing elective major abdominal surgery: a randomized blinded controlled trial. *Ann Surg* 2018;267:50-6.
87. Li C, Carli F, Lee L, Charlebois P, Stein B, Liberman AS, et al. Impact of a trimodal prehabilitation program on functional recovery after colorectal cancer surgery: a pilot study. *Surg Endosc* 2013;27:1072-82.
88. Gillis C, Li C, Lee L, Awasthi R, Augustin B, Gamsa A, et al. Prehabilitation versus rehabilitation: a randomized control trial in patients undergoing colorectal resection for cancer. *Anesthesiology* 2014;121:937-47.
89. Engelman DT, Ali WB, Williams JB, Perrault LP, Reddy VS, Arora RC, et al. Guidelines for perioperative care in cardiac surgery: Enhanced Recovery After Surgery Society recommendations. *JAMA Surg* 2019;154:755-66.
90. Melloul E, Lassen K, Roulin D, Grass F, Perinel J, Adham M, et al. Guidelines for perioperative care for pancreatoduodenectomy: Enhanced Recovery After Surgery (ERAS) recommendations 2019. *World J Surg* 2020;44:2056-84.
91. Wee IJY, Choong A. A systematic review of the impact of preoperative exercise for patients with abdominal aortic aneurysm. *J Vasc Surg* 2020;71:2123-2131.e1.
92. Barakat HM, Shahin Y, Khan JA, McCollum PT, Chetter IC. Preoperative supervised exercise improves outcomes after elective abdominal aortic aneurysm repair: a randomized controlled trial. *Ann Surg* 2016;264:47-53.
93. Dronkers J, Veldman A, Hoberg E, van der Waal C, van Meeteren N. Prevention of pulmonary complications after upper abdominal surgery by preoperative intensive inspiratory muscle training: a randomized controlled pilot study. *Clin Rehabil* 2008;22:134-42.
94. Tew GA, Moss J, Crank H, Mitchell PA, Nawaz S. Endurance exercise training in patients with small abdominal aortic aneurysm: a randomized controlled pilot study. *Arch Phys Med Rehabil* 2012;93:2148-53.

95. Hayashi K, Hirashiki A, Kodama A, Kobayashi K, Yasukawa Y, Shimizu M, et al. Impact of preoperative regular physical activity on postoperative course after open abdominal aortic aneurysm surgery. *Heart Vessels* 2016;31:578-83.
96. Oscarsson A, Gupta A, Fredrikson M, Jarhult J, Nystrom M, Pettersson E, et al. To continue or discontinue aspirin in the perioperative period: a randomized, controlled clinical trial. *Br J Anaesth* 2010;104:305-12.
97. Feringa HHH, Bax JJ, Karagiannis SE, Noordzij P, van Domburg R, Klein J, et al. Elderly patients undergoing major vascular surgery: risk factors and medication associated with risk reduction. *Arch Gerontol Geriatr* 2009;48:116-20.
98. Levine GN, Bates ER, Bittl JA, Brindis RG, Fihn SD, Fleisher LA, et al. 2016 ACC/AHA guideline focused update on duration of dual antiplatelet therapy in patients with coronary artery disease: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines: an update of the 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention, 2011 ACCF/AHA guideline for coronary artery bypass graft surgery, 2012 ACC/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease, 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction, 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes, and 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery. *Circulation* 2016;134:e123-55.
99. Holcomb CN, Graham LA, Richman JS, Rhyne RR, Itani KMF, Maddox TM, et al. The incremental risk of noncardiac surgery on adverse cardiac events following coronary stenting. *J Am Coll Cardiol* 2014;64:2730-9.
100. Holcomb CN, Hollis RH, Graham LA, Richman JS, Valle JA, Itani KM, et al. Association of coronary stent indication with postoperative outcomes following noncardiac surgery. *JAMA Surg* 2016;151:462-9.
101. Cruden NL, Harding SA, Flapan AD, Graham C, Wild SH, Slack R, et al. Previous coronary stent implantation and cardiac events in patients undergoing noncardiac surgery. *Circ Cardiovasc Interv* 2010;3:236-42.
102. Hawn MT, Graham LA, Richman JS, Itani KM, Henderson WG, Maddox TM. Risk of major adverse cardiac events following noncardiac surgery in patients with coronary stents. *JAMA* 2013;310:1462-72.
103. Rossini R, Tarantini G, Musumeci G, Masiero G, Barbato E, Calabro P, et al. A multidisciplinary approach on the perioperative antithrombotic management of patients with coronary stents undergoing surgery: surgery after stenting 2. *JACC Cardiovasc Interv* 2018;11:417-34.
104. Association of Anaesthetists of Great Britain & Ireland; Obstetric Anaesthetists' Association; Regional Anaesthesia UK. Cook T, Gill H, Hill D, et al. Regional anaesthesia and patients with abnormalities of coagulation: the Association of Anaesthetists of Great Britain & Ireland The Obstetric Anaesthetists' Association Regional Anaesthesia UK. *Anaesthesia* 2013;68:966-72.
105. Horlocker TT, Vandermeulen E, Kopp SL, Gogarten W, Leffert LR, Benzon HT. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine evidence-based guidelines (fourth edition). *Reg Anesth Pain Med* 2018;43:263-309.
106. Spahn DR, Munoz M, Klein AA, Levy JH, Zacharowski K. Patient blood management: effectiveness and future potential. *Anesthesiology* 2020;133:212-22.
107. Smith I, Kranke P, Murat I, Smith A, O'Sullivan G, Soreide E, et al. Perioperative fasting in adults and children: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol* 2011;28:556-69.
108. Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures: an updated report by the American Society of Anesthesiologists task force on preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration. *Anesthesiology* 2017;126:376-93.
109. Hausel J, Nygren J, Thorell A, Lagerkranser M, Ljungqvist O. Randomized clinical trial of the effects of oral preoperative carbohydrates on postoperative nausea and vomiting after laparoscopic cholecystectomy. *Br J Surg* 2005;92:415-21.
110. Gianotti L, Biffi R, Sandini M, Marrelli D, Vignali A, Caccialanza R, et al. Preoperative oral carbohydrate load versus placebo in major elective abdominal surgery (PROCY): a randomized, placebo-controlled, multicenter, phase III trial. *Ann Surg* 2018;267:623-30.
111. Wang ZG, Wang Q, Wang WJ, Qin HL. Randomized clinical trial to compare the effects of preoperative oral carbohydrate versus placebo on insulin resistance after colorectal surgery. *Br J Surg* 2010;97:317-27.
112. Faria MS, de Aguiar-Nascimento JE, Pimenta OS, Alvarenga LC Jr, Dock-Nascimento DB, Silhessarenko N. Preoperative fasting of 2 hours minimizes insulin resistance and organic response to trauma after video-cholecystectomy: a randomized, controlled, clinical trial. *World J Surg* 2009;33:1158-64.
113. Rapp-Kesek D, Stridsberg M, Andersson LG, Berne C, Karlsson T. Insulin resistance after cardiopulmonary bypass in the elderly patient. *Scand Cardiovasc J* 2007;41:102-8.
114. Noblett SE, Watson DS, Huong H, Davison B, Hainsworth PJ, Horgan AF. Pre-operative oral carbohydrate loading in colorectal surgery: a randomized controlled trial. *Colorectal Dis* 2006;8:563-9.
115. Lee B, Soh S, Shim JK, Kim HY, Lee H, Kwak YL. Evaluation of preoperative oral carbohydrate administration on insulin resistance in off-pump coronary artery bypass patients: a randomised trial. *Eur J Anaesthesiol* 2017;34:740-7.
116. Feguri GR, Lima PR, Lopes AM, Roledo A, Marchese M, Trevisan M, et al. Clinical and metabolic results of fasting abbreviation with carbohydrates in coronary artery bypass graft surgery. *Rev Bras Cir Cardiovasc* 2012;27:7-17.
117. Breuer JP, von Dossow V, von Heymann C, Griesbach M, von Schickfus M, Mackh E, et al. Preoperative oral carbohydrate administration to ASA III-IV patients undergoing elective cardiac surgery. *Anesth Analg* 2006;103:1099-108.
118. Noba L, Wakefield A. Are carbohydrate drinks more effective than preoperative fasting: a systematic review of randomised controlled trials. *J Clin Nurs* 2019;28:3096-116.
119. Smith MD, McCall J, Plank L, Herbison GP, Soop M, Nygren J. Preoperative carbohydrate treatment for enhancing recovery after elective surgery. *Cochrane Database Syst Rev* 2014;8:CD009161.
120. Gustafsson UO, Nygren J, Thorell A, Soop M, Hellstrom PM, Ljungqvist O, et al. Pre-operative carbohydrate loading may be used in type 2 diabetes patients. *Acta Anaesthesiol Scand* 2008;52:946-51.
121. Laffin MR, Li S, Brisebois R, Senior PA, Wang H. The use of a preoperative carbohydrate drink in patients with diabetes mellitus: a prospective, non-inferiority, cohort study. *World J Surg* 2018;42:1965-70.
122. Farkas JC, Chapuis C, Combe S, Silsiquen M, Marzelle J, Laurian C, et al. A randomised controlled trial of a low-molecular-weight heparin (Enoxaparin) to prevent deep-vein thrombosis in patients undergoing vascular surgery. *Eur J Vasc Surg* 1993;7:554-60.
123. Pawlaczyk K, Gabriel M, Dzieciuchowicz L, Stanisic M, Begier-Krasinska B, Gabriel Z, et al. Post-operative venous thromboembolism in patients operated on for aorto-iliac obstruction and abdominal aortic aneurysm, and the application of pharmacological thromboprophylaxis. *Eur J Vasc Endovasc Surg* 2016;51:121-6.
124. de Maistre E, Terriat B, Lesne-Padieu AS, Abello N, Bouchot O, Steinmetz EF. High incidence of venous thrombosis after surgery for abdominal aortic aneurysm. *J Vasc Surg* 2009;49:596-601.
125. Khan NK, Oksala NK, Suominen V, Vakhitov D, JO Laurikka, Khan JA. Risk of symptomatic venous thromboembolism after abdominal aortic aneurysm repair in long-term follow-up of 1021 consecutive patients. *J Vasc Surg Venous Lymphat Disord* 2021;9:54-61.
126. Ramanan B, Gupta PK, Sundaram A, Lynch TG, MacTaggart JN, Baxter BT, et al. In-hospital and postdischarge venous thromboembolism after vascular surgery. *J Vasc Surg* 2013;57:1589-96.
127. Toth S, Flohr TR, Schubart J, Knehans A, Castello MC, Aziz F. A meta-analysis and systematic review of venous thromboembolism prophylaxis in patients undergoing vascular surgery procedures. *J Vasc Surg Venous Lymphat Disord* 2020;8:869-81.e2.
128. Killewich LA, Aswad MA, Sandager GP, Lilly MP, Flinn WR. A randomized, prospective trial of deep venous thrombosis prophylaxis in aortic surgery. *Arch Surg* 1997;132:499-504.

129. Gould MK, Garcia DA, Wren SM, Karanicolas PJ, Arcelus JI, Heit JA, et al. Prevention of VTE in nonorthopedic surgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2012;141(Suppl):e227S-77S.
130. Nicolaidis AN, Fareed J, Kakkar AK, Comerota AJ, Goldhaber SZ, Hull R, et al. Prevention and treatment of venous thromboembolism—international consensus statement. *Int Angiol* 2013;32:111-260.
131. Ip HY, Abrishami A, Peng PW, Wong J, Chung F. Predictors of postoperative pain and analgesic consumption: a qualitative systematic review. *Anesthesiology* 2009;111:657-77.
132. American Geriatrics Society 2015 Beers Criteria Update Expert Panel. American Geriatrics Society 2015 updated Beers criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc* 2015;63:2227-46.
133. Gaulton TG, Wunsch H, Gaskins LJ, Leonard CE, Hennessy S, Ashburn M, et al. Preoperative sedative-hypnotic medication use and adverse postoperative outcomes. *Ann Surg* 2021;274:e108-14.
134. Wongkietkachorn A, Wongkietkachorn N, Rhunsiri P. Preoperative needs-based education to reduce anxiety, increase satisfaction, and decrease time spent in day surgery: a randomized controlled trial. *World J Surg* 2018;42:666-74.
135. Gustafsson UO, Scott MJ, Schwenk W, Demartines N, Roulin D, Francis N, et al. Guidelines for perioperative care in elective colonic surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *World J Surg* 2013;37:259-84.
136. Torrance N, Veluchamy A, Zhou Y, Fletcher EH, Moir E, Hebert HL, et al. Trends in gabapentinoid prescribing, co-prescribing of opioids and benzodiazepines, and associated deaths in Scotland. *Br J Anaesth* 2020;125:159-67.
137. McAnally H, Bonnet U, Kaye AD. Gabapentinoid benefit and risk stratification: mechanisms over myth. *Pain Ther* 2020;9:441-52.
138. Sheth KR, Bernthal NM, Ho HS, Bergese SD, Apfel CC, Stoicea N, et al. Perioperative bleeding and non-steroidal anti-inflammatory drugs: an evidence-based literature review, and current clinical appraisal. *Medicine (Baltimore)* 2020;99:e20042.
139. Stewart A, Evers PS, Earnshaw JJ. Prevention of infection in arterial reconstruction. *Cochrane Database Syst Rev* 2006;3:CD003073.
140. Engelman R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, et al. The Society of Thoracic Surgeons practice guideline series: antibiotic prophylaxis in cardiac surgery, part II: antibiotic choice. *Ann Thorac Surg* 2007;83:1569-76.
141. Edwards FH, Engelman RM, Houck P, Shahian DM, Bridges CR; Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: antibiotic prophylaxis in cardiac surgery, part I: duration. *Ann Thorac Surg* 2006;81:397-404.
142. Zanetti G, Giardina R, Platt R. Intraoperative redosing of cefazolin and risk for surgical site infection in cardiac surgery. *Emerg Infect Dis* 2001;7:828-31.
143. Bratzler DW, Houck PM. Antimicrobial prophylaxis for surgery: an advisory statement from the National Surgical Infection Prevention Project. *Clin Infect Dis* 2004;38:1706-15.
144. Swoboda SM, Merz C, Kostuik J, Trentler B, Lipsett PA. Does intraoperative blood loss affect antibiotic serum and tissue concentrations? *Arch Surg* 1996;131:1165-71; discussion: 1171, 1162.
145. Gan TJ, Belani KG, Bergese S, Chung F, Diemunsch P, Habib AS, et al. Fourth consensus guidelines for the management of postoperative nausea and vomiting. *Anesth Analg* 2020;131:411-48.
146. Apfel CC, Heidrich FM, Jukar-Rao S, Jalota L, Hornuss C, Whelan RP, et al. Evidence-based analysis of risk factors for postoperative nausea and vomiting. *Br J Anaesth* 2012;109:742-53.
147. Apfel CC, Laara E, Koivuranta M, Greim CA, Roewer N. A simplified risk score for predicting postoperative nausea and vomiting: conclusions from cross-validations between two centers. *Anesthesiology* 1999;91:693-700.
148. Koivuranta M, Laara E, Snare L, Alahuhta S. A survey of postoperative nausea and vomiting. *Anaesthesia* 1997;52:443-9.
149. Apfel CC, Korttila K, Abdalla M, Kerger H, Turan A, Vedder I, et al. A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *N Engl J Med* 2004;350:2441-51.
150. Liu SS, Strödtbeck WM, Richman JM, Wu CL. A comparison of regional versus general anesthesia for ambulatory anesthesia: a meta-analysis of randomized controlled trials. *Anesth Analg* 2005;101:1634-42.
151. Apfel CC, Turan A, Souza K, Pergolizzi J, Hornuss C. Intravenous acetaminophen reduces postoperative nausea and vomiting: a systematic review and meta-analysis. *Pain* 2013;154:677-89.
152. Elia N, Lysakowski C, Tramer MR. Does multimodal analgesia with acetaminophen, nonsteroidal antiinflammatory drugs, or selective cyclooxygenase-2 inhibitors and patient-controlled analgesia morphine offer advantages over morphine alone? Meta-analyses of randomized trials. *Anesthesiology* 2005;103:1296-304.
153. Cao X, White PF, Ma H. An update on the management of postoperative nausea and vomiting. *J Anesth* 2017;31:617-26.
154. Kaye AD, Cornett EM, Chalabi J, Naim NZ, Novitch MB, Creel JB, et al. Pharmacology of antiemetics: update and current considerations in anesthesia practice. *Anesthesiol Clin* 2017;35:e41-54.
155. McLean DJ, Diaz-Gil D, Farhan HN, Ladha KS, Kurth T, Eikermann M. Dose-dependent association between intermediate-acting neuromuscular-blocking agents and postoperative respiratory complications. *Anesthesiology* 2015;122:1201-13.
156. Brull SJ, Murphy GS. Residual neuromuscular block: lessons unlearned. Part II: methods to reduce the risk of residual weakness. *Anesth Analg* 2010;111:129-40.
157. Brull SJ, Kopman AF. Current status of neuromuscular reversal and monitoring: challenges and opportunities. *Anesthesiology* 2017;126:173-90.
158. Kheterpal S, Vaughn MT, Dubovoy TZ, Shah NJ, Bash LD, Colquhoun DA, et al. Sugammadex versus neostigmine for reversal of neuromuscular blockade and postoperative pulmonary complications (STRONGER): a multicenter matched cohort analysis. *Anesthesiology* 2020;132:1371-81.
159. Oh TK, Oh A-Y, Ryu J-H, Koo B-W, Song I-A, Nam SW, et al. Retrospective analysis of 30-day unplanned readmission after major abdominal surgery with reversal by sugammadex or neostigmine. *Br J Anaesth* 2019;122:370-8.
160. Togioka BM, Yanez D, Aziz MF, Higgins JR, Tekkali P, Treggiari MM. Randomised controlled trial of sugammadex or neostigmine for reversal of neuromuscular block on the incidence of pulmonary complications in older adults undergoing prolonged surgery. *Br J Anaesth* 2020;124:553-61.
161. Bender JS, Smith-Meek MA, Jones CE. Routine pulmonary artery catheterization does not reduce morbidity and mortality of elective vascular surgery: results of a prospective, randomized trial. *Ann Surg* 1997;226:229-36; discussion: 236.
162. Bonazzi M, Gentile F, Biasi GM, Migliavacca S, Esposti D, Cipolla M, et al. Impact of perioperative haemodynamic monitoring on cardiac morbidity after major vascular surgery in low risk patients: a randomised pilot trial. *Eur J Vasc Endovasc Surg* 2002;23:445-51.
163. American Society of Anesthesiologists; Society of Cardiovascular Anesthesiologists Task Force on Transesophageal Echocardiography. Practice guidelines for perioperative transesophageal echocardiography: an updated report by the American Society of Anesthesiologists and the Society of Cardiovascular Anesthesiologists Task Force on Transesophageal Echocardiography. *Anesthesiology* 2010;112:1084-96.
164. Fayad A, Shillcutt SK. Perioperative transesophageal echocardiography for non-cardiac surgery. *Can J Anaesth* 2018;65:381-98.
165. Hadian M, Pinsky MR. Evidence-based review of the use of the pulmonary artery catheter: impact data and complications. *Crit Care* 2006;10(Suppl 3):S8.
166. Walsh M, Devereaux PJ, Garg AX, Kurz A, Turan A, Rodseth RN, et al. Relationship between intraoperative mean arterial pressure and clinical outcomes after noncardiac surgery: toward an empirical definition of hypotension. *Anesthesiology* 2013;119:507-15.
167. van Waas JAR, van Klei WA, Wijjensundera DN, van Wolfswinkel L, Lindsay TF, Beattie WS. Association between intraoperative hypotension and myocardial injury after vascular surgery. *Anesthesiology* 2016;124:35-44.
168. Salmasi V, Maheshwari K, Yang D, Mascha EJ, Singh A, Sessler DI, et al. Relationship between intraoperative hypotension, defined by either reduction from baseline or absolute thresholds, and acute kidney and myocardial injury after noncardiac surgery: a retrospective cohort analysis. *Anesthesiology* 2017;126:47-65.

169. National Institute for Health Care Excellence. Abdominal aortic aneurysm: diagnosis and management. Available at: <https://www.nice.org.uk/guidance/ng156>. Accessed December 10, 2020.
170. De Freitas S, Hicks CW, Mouton R, Garcia S, Healy D, Connolly C, et al. Effects of ischemic preconditioning on abdominal aortic aneurysm repair: a systematic review and meta-analysis. *J Surg Res* 2019;235:340-9.
171. Baldwin L, Henderson A, Hickman P. Effect of postoperative low-dose dopamine on renal function after elective major vascular surgery. *Ann Intern Med* 1994;120:744-7.
172. Gillies MA, Kakar V, Parker RJ, Honore PM, Ostermann M. Fenoldopam to prevent acute kidney injury after major surgery—a systematic review and meta-analysis. *Crit Care* 2015;19:449.
173. Halpenny M, Rushe C, Breen P, Cunningham AJ, Boucher-Hayes D, Shorten GD. The effects of fenoldopam on renal function in patients undergoing elective aortic surgery. *Eur J Anaesthesiol* 2002;19:32-9.
174. Khwaja A. KDIGO clinical practice guidelines for acute kidney injury. *Nephron Clin Pract* 2012;120:c179-84.
175. Hersey P, Poullis M. Does the administration of mannitol prevent renal failure in open abdominal aortic aneurysm surgery? *Interact Cardiovasc Thorac Surg* 2008;7:906-9.
176. Dubois L, Durant C, Harrington DM, Forbes TL, Derose G, Harris JR. Technical factors are strongest predictors of postoperative renal dysfunction after open transperitoneal juxtarenal abdominal aortic aneurysm repair. *J Vasc Surg* 2013;57:648-54.
177. Deery SE, Lancaster RT, Baril DT, Indes JE, Bertges DJ, Conrad MF, et al. Contemporary outcomes of open complex abdominal aortic aneurysm repair. *J Vasc Surg* 2016;63:1195-200.
178. Shantikumar S, Patel S, Handa A. The role of cell salvage autotransfusion in abdominal aortic aneurysm surgery. *Eur J Vasc Endovasc Surg* 2011;42:577-84.
179. Takagi H, Sekino S, Kato T, Matsuno Y, Umemoto T. Intraoperative autotransfusion in abdominal aortic aneurysm surgery: meta-analysis of randomized controlled trials. *Arch Surg* 2007;142:1098-101.
180. Klein AA, Bailey CR, Charlton AJ, Evans E, Guckian-Fisher M, McCrossan R, et al. Association of Anaesthetists guidelines: cell salvage for peri-operative blood conservation 2018. *Anaesthesia* 2018;73:1141-50.
181. Odor PM, Bampoe S, Gilhooly D, Creagh-Brown B, Moonesinghe SR. Perioperative interventions for prevention of postoperative pulmonary complications: systematic review and meta-analysis. *BMJ* 2020;368:m540.
182. Futier E, Constantin J-M, Paugam-Burtz C, Pascal J, Eurin M, Neuschwander A, et al. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med* 2013;369:428-37.
183. Culdner A, Kiss T, Serpa Neto A, Hemmes SNT, Canet J, Spieth PM, et al. Intraoperative protective mechanical ventilation for prevention of postoperative pulmonary complications: a comprehensive review of the role of tidal volume, positive end-expiratory pressure, and lung recruitment maneuvers. *Anesthesiology* 2015;123:692-713.
184. Park WY, Thompson JS, Lee KK. Effect of epidural anesthesia and analgesia on perioperative outcome: a randomized, controlled Veterans Affairs cooperative study. *Ann Surg* 2001;234:560-9; discussion: 569-71.
185. Peyton PJ, Myles PS, Silbert BS, Rigg JA, Jamrozik K, Parsons R. Perioperative epidural analgesia and outcome after major abdominal surgery in high-risk patients. *Anesth Analg* 2003;96:548. table of contents.
186. Broekema AA, Veen A, Fidler V, Gielen MJ, Hennis PJ. Postoperative analgesia with intramuscular morphine at fixed rate versus epidural morphine or sufentanil and bupivacaine in patients undergoing major abdominal surgery. *Anesth Analg* 1998;87:1346-53.
187. Boylan JF, Katz J, Kavanagh BP, Klinck JR, Cheng DC, DeMajo WC, et al. Epidural bupivacaine-morphine analgesia versus patient-controlled analgesia following abdominal aortic surgery: analgesic, respiratory, and myocardial effects. *Anesthesiology* 1998;89:585-93.
188. Bois S, Couture P, Boudreault D, Lacombe P, Fugere F, Girard D, et al. Epidural analgesia and intravenous patient-controlled analgesia result in similar rates of postoperative myocardial ischemia after aortic surgery. *Anesth Analg* 1997;85:1233-9.
189. Renghi A, Gramaglia L, Casella F, Moniaci D, Gaboli K, Brustia P. Local versus epidural anesthesia in fast-track abdominal aortic surgery. *J Cardiothorac Vasc Anesth* 2013;27:451-8.
190. Norris EJ, Beattie C, Perler BA, Martinez EA, Meinert CL, Anderson GF, et al. Double-masked randomized trial comparing alternate combinations of intraoperative anesthesia and postoperative analgesia in abdominal aortic surgery. *Anesthesiology* 2001;95:1054-67.
191. Guay J, Kopp S. Epidural pain relief versus systemic opioid-based pain relief for abdominal aortic surgery. *Cochrane Database Syst Rev* 2016;1:CD005059.
192. National Institutes of Health and Care Excellence. Hypothermia: prevention and management in adults having surgery. NICE Clinical Guidelines, No. 65. Available at: <https://www.nice.org.uk/guidance/cg65>. Accessed December 10, 2020.
193. Healy D, Clarke-Moloney M, Hannigan A, Walsh S. Wound drainage for lower limb arterial surgery. *Cochrane Database Syst Rev* 2016;11:CD011111.
194. Kwon J, Staley C, McCullough M, Goss S, Arosemena M, Abai B, et al. A randomized clinical trial evaluating negative pressure therapy to decrease vascular groin incision complications. *J Vasc Surg* 2018;68:1744-52.
195. Matatov T, Reddy KN, Doucet LD, Zhao CX, Zhang WW. Experience with a new negative pressure incision management system in prevention of groin wound infection in vascular surgery patients. *J Vasc Surg* 2013;57:791-5.
196. Lee K, Murphy PB, Ingves MV, Duncan A, DeRose G, Dubois L, et al. Randomized clinical trial of negative pressure wound therapy for high-risk groin wounds in lower extremity revascularization. *J Vasc Surg* 2017;66:1814-9.
197. Popping DM, Elia N, Van Aken HK, Marret E, Schug SA, Kranke P, et al. Impact of epidural analgesia on mortality and morbidity after surgery: systematic review and meta-analysis of randomized controlled trials. *Ann Surg* 2014;259:1056-67.
198. Brown EN, Pavone KJ, Naranjo M. Multimodal general anesthesia: theory and practice. *Anesth Analg* 2018;127:1246-58.
199. Cerantola Y, Valerio M, Persson B, Jichlinski P, Ljungqvist O, Hubner M, et al. Guidelines for perioperative care after radical cystectomy for bladder cancer: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *Clin Nutr* 2013;32:879-87.
200. Scott MJ, Baldini G, Fearon KC, Feldheiser A, Feldman LS, Gan TJ, et al. Enhanced Recovery After Surgery (ERAS) for gastrointestinal surgery, part 1: pathophysiological considerations. *Acta Anaesthesiol Scand* 2015;59:1212-31.
201. Kang SH, Lee Y, Min S-H, Park YS, Ahn S-H, Park DJ, et al. Multimodal Enhanced Recovery After Surgery (ERAS) program is the optimal perioperative care in patients undergoing totally laparoscopic distal gastrectomy for gastric cancer: a prospective, randomized, clinical trial. *Ann Surg Oncol* 2018;25:3231-8.
202. Kumar K, Kirksey MA, Duong S, Wu CL. A review of opioid-sparing modalities in perioperative pain management: methods to decrease opioid use postoperatively. *Anesth Analg* 2017;125:1749-60.
203. Bell RF, Dahl JB, Moore RA, Kalso E. Perioperative ketamine for acute postoperative pain. *Cochrane Database Syst Rev* 2006;1:CD004603.
204. Cheung CC, Martyn A, Campbell N, Frost S, Gilbert K, Michota F, et al. Predictors of intraoperative hypotension and bradycardia. *Am J Med* 2015;128:532-8.
205. Eipe N, Penning J, Yazdi F, Mallick R, Turner L, Ahmadzai N, et al. Perioperative use of pregabalin for acute pain—a systematic review and meta-analysis. *Pain* 2015;156:1284-300.
206. Ge DJ, Qi B, Tang C, Li JY. Intraoperative dexmedetomidine promotes postoperative analgesia and recovery in patients after abdominal hysterectomy: a double-blind, randomized clinical trial. *Sci Rep* 2016;6:21514.
207. Kranke P, Jokinen J, Pace NL, Schnabel A, Hollmann MW, Hahnenkamp K, et al. Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery. *Cochrane Database Syst Rev* 2015;7:CD009642.
208. Bahr MP, Williams BA. Esmolol, antinociception, and its potential opioid-sparing role in routine anesthesia care. *Reg Anesth Pain Med* 2018;43:815-8.

209. Chidgey BA, McGinigle KL, McNaull PP. When a vital sign leads a country astray—the opioid epidemic. *JAMA Surg* 2019;154:987-8.
210. Santosa KB, Hu H-M, Brummett CM, Olsen MA, Englesbe MJ, Williams EA, et al. New persistent opioid use among older patients following surgery: a Medicare claims analysis. *Surgery* 2020;167:732-42.
211. Clegg A, Young JB. Which medications to avoid in people at risk of delirium: a systematic review. *Age Ageing* 2011;40:23-9.
212. Duprey MS, Dijkstra-Kersten SMA, Zaal IJ, Briesacher BA, Saczynski JS, Griffith JL, et al. Opioid use increases the risk of delirium in critically ill adults independently of pain. *Am J Respir Crit Care Med* 2021;204:566-72.
213. Benzon HT, Raja S, Liu SS, Fishman S, Cohen SP. *Essentials of Pain Medicine*. 4th ed. 2018. Elsevier.
214. Wei ZW, Li JL, Li ZS, Hao YT, He YL, Chen W, et al. Systematic review of nasogastric or nasojejunal decompression after gastrectomy for gastric cancer. *Eur J Surg Oncol* 2014;40:1763-70.
215. Wang D, Li T, Yu J, Hu Y, Liu H, Li G. Is nasogastric or nasojejunal decompression necessary following gastrectomy for gastric cancer? A systematic review and meta-analysis of randomised controlled trials. *J Gastrointest Surg* 2015;19:195-204.
216. Vermeulen H, Storm-Versloot MN, Busch OR, Ubbink DT. Nasogastric intubation after abdominal surgery: a meta-analysis of recent literature. *Arch Surg* 2006;141:307-14.
217. Nelson R, Edwards S, Tse B. Prophylactic nasogastric decompression after abdominal surgery. *Cochrane Database Syst Rev* 2007;3:CD004929.
218. Avrahami R, Cohen JD, Haddad M, Singer P, Zelikovski A. Gastric emptying after elective abdominal aortic aneurysm surgery: the case for early postoperative enteral feeding. *Eur J Vasc Endovasc Surg* 1999;17:241-4.
219. Ksienski MR, Fenton TR, Eliasziw M, Zuege DJ, Petrasko P, Shahpori R, et al. A cohort study of nutrition practices in the intensive care unit following abdominal aortic aneurysm repair. *J Parenter Ther* 2013;37:261-7.
220. Martelli M, Renghi A, Gramaglia L, Casella F, Brustia P. Abdominal aortic aneurysm treatment: minimally invasive fast-track surgery and endovascular technique in octogenarians. *J Cardiovasc Surg (Torino)* 2017;58:557-64.
221. Lohr G, Keller H, Kutscher J, Huber R. [Fast-track rehabilitation after open aorto-iliac vascular procedures]. *Zentralbl Chir* 2008;133:344-8.
222. Dass B, Shimada M, Kambhampati G, Ejaz NI, Arif AA, Ejaz AA. Fluid balance as an early indicator of acute kidney injury in CV surgery. *Clin Nephrol* 2012;77:438-44.
223. Myles PS, Bellomo R, Corcoran T, Forbes A, Peyton P, Story D, et al. Restrictive versus liberal fluid therapy for major abdominal surgery. *N Engl J Med* 2018;378:2263-74.
224. Jonsson K, Jensen JA, Goodson WH III, Scheuenstuhl H, West J, Hopf HW, et al. Tissue oxygenation, anemia, and perfusion in relation to wound healing in surgical patients. *Ann Surg* 1991;214:605-13.
225. Thiele RH, Rea KM, Turrentine FE, Friel CM, Hassinger TE, McMurry TL, et al. Standardization of care: impact of an enhanced recovery protocol on length of stay, complications, and direct costs after colorectal surgery. *J Am Coll Surg* 2015;220:430-43.
226. Makaryus R, Miller TE, Gan TJ. Current concepts of fluid management in enhanced recovery pathways. *Br J Anaesth* 2018;120:376-83.
227. Kim Y, Gani F, Spolverato G, Ejaz A, Xu L, Buettner S, et al. Variation in crystalloid administration: an analysis of 6248 patients undergoing major elective surgery. *J Surg Res* 2016;203:368-77.
228. Lilot M, Ehrenfeld JM, Lee C, Harrington B, Cannesson M, Rinehart J. Variability in practice and factors predictive of total crystalloid administration during abdominal surgery: retrospective two-centre analysis. *Br J Anaesth* 2015;114:767-76.
229. Gonzalez-Fajardo JA, Mengibar L, Brizuela JA, Castrodeza J, Vaquero-Puerta C. Effect of postoperative restrictive fluid therapy in the recovery of patients with abdominal vascular surgery. *Eur J Vasc Endovasc Surg* 2009;37:538-43.
230. McArdle GT, McAuley DF, McKinley A, Blair P, Hoper M, Harkin DW. Preliminary results of a prospective randomized trial of restrictive versus standard fluid regime in elective open abdominal aortic aneurysm repair. *Ann Surg* 2009;250:28-34.
231. Piljic D, Petricevic M, Piljic D, Ksela J, Robic B, Klokocovnik T. Restrictive versus standard fluid regimen in elective minilaparotomy abdominal aortic repair—prospective randomized controlled trial. *Thorac Cardiovasc Surg* 2016;64:296-303.
232. Toomtong P, Suksompong S. Intravenous fluids for abdominal aortic surgery. *Cochrane Database Syst Rev* 2010;1:CD000991.
233. Wald R, Waikar SS, Liangos O, Pereira BJC, Chertow GM, Jaber BL. Acute renal failure after endovascular vs open repair of abdominal aortic aneurysm. *J Vasc Surg* 2006;43:460-6; discussion: 466.
234. Nonaka T, Kimura N, Hori D, Sasabuchi Y, Nakano M, Yuri K, et al. Predictors of acute kidney injury following elective open and endovascular aortic repair for abdominal aortic aneurysm. *Ann Vasc Dis* 2018;11:298-305.
235. Dang T, Dakour-Aridi H, Rizwan M, Nejm B, Malas MB. Predictors of acute kidney injury after infrarenal abdominal aortic aneurysm repair in octogenarians. *J Vasc Surg* 2019;69:752-62.e1.
236. Gould CV, Umscheid CA, Agarwal RK, Kuntz C, Pegues DA. Healthcare Infection Control Practices Advisory Committee. Guideline for prevention of catheter-associated urinary tract infections 2009. *Infect Control Hosp Epidemiol* 2010;31:319-26.
237. Ladak SS, Katznelson R, Muscat M, Sawhney M, Beattie WS, O'Leary G. Incidence of urinary retention in patients with thoracic patient-controlled epidural analgesia (TPCEA) undergoing thoracotomy. *Pain Manag Nurs* 2009;10:94-8.
238. De Leon LE, Patil N, Hartigan PM, White A, Bravo-Iniguez CE, Fox S, et al. Risk of urinary recatheterization for thoracic surgical patients with epidural anesthesia. *J Surg Res (Houst)* 2020;3:163-71.
239. Subramaniam B, Panzica PJ, Novack V, Mahmood F, Matyal R, Mitchell JD, et al. Continuous perioperative insulin infusion decreases major cardiovascular events in patients undergoing vascular surgery: a prospective, randomized trial. *Anesthesiology* 2009;110:970-7.
240. Long CA, Fang ZB, Hu FY, Arya S, Brewster LP, Duggan E, et al. Poor glycemic control is a strong predictor of postoperative morbidity and mortality in patients undergoing vascular surgery. *J Vasc Surg* 2019;69:1219-26.
241. Action to Control Cardiovascular Risk in Diabetes Study Group, Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, et al. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545-59.
242. NICE-SUGAR Study Investigators, Finfer S, Liu B, Chittock DR, Norton R, Myburgh JA, et al. Hypoglycemia and risk of death in critically ill patients. *N Engl J Med* 2012;367:1108-18.
243. Duwayri Y, Jordan WD Jr. Diabetes, dysglycemia, and vascular surgery. *J Vasc Surg* 2020;71:701-11.
244. Stethen TW, Ghazi YA, Heidel RE, Daley BJ, Barnes L, Patterson D, et al. Walking to recovery: the effects of missed ambulation events on postsurgical recovery after bowel resection. *J Gastrointest Oncol* 2018;9:953-61.
245. Kalisch BJ, Lee S, Dabney BW. Outcomes of inpatient mobilization: a literature review. *J Clin Nurs* 2014;23:1486-501.
246. Shallom M, Tymkew H, Vyers K, Prentice D, Sona C, Norris T, et al. Implementation of an interdisciplinary AACN early mobility protocol. *Crit Care Nurse* 2020;40:e7-17.
247. Fiore JF Jr, Bialocerkowski A, Browning L, Faragher IG, Denehy L. Criteria to determine readiness for hospital discharge following colorectal surgery: an international consensus using the Delphi technique. *Dis Colon Rectum* 2012;55:416-23.
248. Rosen SF, Clagett GP, Valentine RJ, Jackson MR, Modrall JC, McIntyre KE. Transient advanced mental impairment: an underappreciated morbidity after aortic surgery. *J Vasc Surg* 2002;35:376-81.
249. Tahhan G, Farber A, Shah NK, Krafcik BM, Sachs TE, Kalish JA, et al. Characterization of planned and unplanned 30-day readmissions following vascular surgical procedures. *Vasc Endovascular Surg* 2017;51:17-22.
250. Knighton A, Martin G, Sounderajah V, Warren L, Markiewicz O, Riga C, et al. Avoidable 30-day readmissions in patients undergoing vascular surgery. *BJS Open* 2019;3:759-66.
251. Eun JC, Nehler MR, Black JH III, Glebova NO. Measures to reduce unplanned readmissions after vascular surgery. *Semin Vasc Surg* 2015;28:103-11.
252. Brooke BS, De Martino RR, Girotti M, Dimick JB, Goodney PP. Developing strategies for predicting and preventing readmissions in vascular surgery. *J Vasc Surg* 2012;56:556-62.

253. Johnson A, Sandford J, Tyndall J. Written and verbal information versus verbal information only for patients being discharged from acute hospital settings to home. *Cochrane Database Syst Rev* 2003;4:CD003716.
254. Li LT, Mills WL, Gutierrez AM, Herman LI, Berger DH, Naik AD. A patient-centered early warning system to prevent readmission after colorectal surgery: a national consensus using the Delphi method. *J Am Coll Surg* 2013;216:210-6.e6.
255. Mistiaen P, Poot E. Telephone follow-up, initiated by a hospital-based health professional, for postdischarge problems in patients discharged from hospital to home. *Cochrane Database Syst Rev* 2006;4:CD004510.
256. Brooke BS, Stone DH, Cronenwett JL, Nolan B, DeMartino RR, MacKenzie TA, et al. Early primary care provider follow-up and readmission after high-risk surgery. *JAMA Surg* 2014;149:821-8.
257. Ripolles-Melchor J, Ramirez-Rodriguez JM, Casans-Frances R, Aldecoa C, Abad-Motos A, Logrono-Egea M, et al. Association between use of enhanced recovery after surgery protocol and postoperative complications in colorectal surgery: the postoperative outcomes within enhanced recovery after surgery protocol (PO-WER) study. *JAMA Surg* 2019;154:725-36.
258. Currie A, Soop M, Demartines N, Fearon K, Kennedy R, Ljungqvist O. Enhanced recovery after surgery interactive audit system: 10 years' experience with an international web-based clinical and research perioperative care database. *Clin Colon Rectal Surg* 2019;32:75-81.
259. Gustafsson UO, Hausel J, Thorell A, Ljungqvist O, Soop M, Nygren J, et al. Adherence to the enhanced recovery after surgery protocol and outcomes after colorectal cancer surgery. *Arch Surg* 2011;146:571-7.
260. Bisch SP, Wells T, Gramlich L, Faris P, Wang X, Tran DT, et al. Enhanced Recovery After Surgery (ERAS) in gynecologic oncology: system-wide implementation and audit leads to improved value and patient outcomes. *Gynecol Oncol* 2018;151:117-23.
261. de Groot JJA, van Es LEJM, Maessen JMC, Dejong CHC, Kruitwagen RFFM, Slangen BFM. Diffusion of enhanced recovery principles in gynecologic oncology surgery: is active implementation still necessary? *Gynecol Oncol* 2014;134:570-5.
262. Cakir H, van Stijn MF, Lopes Cardozo AM, Langenhorst BL, Schreurs WH, van der Ploeg TJ, et al. Adherence to enhanced recovery after surgery and length of stay after colonic resection. *Colorectal Dis* 2013;15:1019-25.
263. Committee on Standards and Practice Parameters, Apfelbaum JL, Connis RT, Nickinovich DG. American Society of Anesthesiologists Task Force on Preanesthesia Evaluation; Pasternak LR, et al. Practice advisory for preanesthesia evaluation: an updated report by the American Society of Anesthesiologists task force on preanesthesia evaluation. *Anesthesiology* 2012;116:522-38.
264. Qaseem A, Snow V, Fitterman N, Hornbake ER, Lawrence VA, Smetana GW, et al. Risk assessment for and strategies to reduce perioperative pulmonary complications for patients undergoing noncardiothoracic surgery: a guideline from the American College of Physicians. *Ann Intern Med* 2006;144:575-80.
265. Expert Panel on Thoracic Imaging, McComb BL, Chung JH, Crabtree TD, Heitkamp DE, Iannettoni MD, et al. ACR Appropriateness Criteria® routine chest radiography. *J Thorac Imaging* 2016;31:W13-5.
266. Yokota S, Koizumi M, Togashi K, Morimoto M, Yasuda Y, Sata N, et al. Preoperative pulmonary function tests do not predict the development of pulmonary complications after elective major abdominal surgery: a prospective cohort study. *Int J Surg* 2020;73:65-71.
267. American Society of Anesthesiologists Task Force on Preanesthesia Evaluation. Practice advisory for preanesthesia evaluation: a report by the American Society of Anesthesiologists Task Force on Preanesthesia Evaluation. *Anesthesiology* 2002;96:485-96.
268. National Institute for Health and Care Excellence. Routine preoperative tests for elective surgery. Available at: <https://www.nice.org.uk/guidance/ng45>. Accessed December 10, 2020.
269. Ho P, Ting AC, Cheng SW. Blood loss and transfusion in elective abdominal aortic aneurysm surgery. *ANZ J Surg* 2004;74:631-4.
270. Matsumura JS, Katzen BT, Sullivan TM, Dake MD, Naftel DC; Excluder Bifurcated Endoprosthesis Investigators. Predictors of survival following open and endovascular repair of abdominal aortic aneurysms. *Ann Vasc Surg* 2009;23:153-8.
271. van Hendriks Y, Romme J, der Slegt J, van Veen EJ, der Laan L, Kluytmans J. Implementation of a bundle of care to reduce surgical site infections in patients undergoing vascular surgery. *PLoS One* 2013;8:e71566.
272. Tanner J, Norrie P, Melen K. Preoperative hair removal to reduce surgical site infection. *Cochrane Database Syst Rev* 2011;11:CD004122.
273. Leaper D, Burman-Roy S, Palanca A, Cullen K, Worster D, Gautam-Aitken E, et al. Prevention and treatment of surgical site infection: summary of NICE guidance. *BMJ* 2008;337:a1924.
274. Darouiche RO, Wall MJ Jr, Itani KM, Otterson MF, Webb AL, Carrick MM, et al. Chlorhexidine-alcohol versus povidone-iodine for surgical-site antisepsis. *N Engl J Med* 2010;362:18-26.
275. Earnshaw JJ, Berridge DC, Slack RC, Makin GS, Hopkinson BR. Do preoperative chlorhexidine baths reduce the risk of infection after vascular reconstruction? *Eur J Vasc Surg* 1989;3:323-6.
276. Fernandez-Prada M, Martinez-Ortega C, Revuelta-Marino L, Menendez-Herrero A, Navarro-Gracia JF. Evaluation of the bundle "zero surgical site infection" to prevent surgical site infection in vascular surgery. *Ann Vasc Surg* 2017;41:160-8.
277. Ljungqvist O, de Boer HD, Balfour A, Fawcett WJ, Lobo DN, Nelson G, et al. Opportunities and challenges for the next phase of enhanced recovery after surgery: a review. *JAMA Surg* 2021;156:775-84.

Submitted Dec 15, 2021; accepted Jan 3, 2022.